


Team announces breakthrough observation of Mott transition in a superconductor

11 September 2015, by Joost Bruysters



quantum or classical, but not both.

One system that unites the two is found in superconductors, certain materials that conduct electricity perfectly when cooled to very low temperatures. Magnetic fields penetrate the superconducting material in the form of tiny filaments called vortices, which control the electronic and magnetic properties of the materials.

These vortices display both classical and quantum properties, which led researchers to study them for access to one of the most enigmatic phenomena of modern [condensed matter physics](#): the Mott insulator-to-metal transition.


An international team of researchers, including the MESA+ Institute for Nanotechnology at the University of Twente in The Netherlands and the U.S. Department of Energy's Argonne National Laboratory, announced today in *Science* the observation of a dynamic Mott transition in a superconductor.

The discovery experimentally connects the worlds of classical and quantum mechanics and illuminates the mysterious nature of the Mott transition. It also could shed light on non-equilibrium [physics](#), which is poorly understood but governs most of what occurs in our world. The finding may also represent a step towards more efficient electronics based on the Mott transition.

Since its foundations were laid in the early part of the 20th century, scientists have been trying to reconcile quantum mechanics with the rules of classical or Newtonian physics (like how you describe the path of an apple thrown into the air—or dropped from a tree). Physicists have made strides in linking the two approaches, but experiments that connect the two are still few and far between; physics phenomena are usually classified as either


The Mott transition occurs in certain materials that according to textbook [quantum mechanics](#) should be metals, but in reality turn insulators. A complex phenomenon controlled by the interactions of many quantum particles, the Mott transition remains mysterious—even whether or not it's a classical or quantum phenomenon is not quite clear. Moreover, scientists have never directly observed a dynamic Mott transition, in which a phase transition from an insulating to a metallic state is induced by driving an electrical current through the system; the disorder inherent in real systems disguises Mott properties.

At the University of Twente, researchers built a system containing 90,000 superconducting niobium nano-sized islands on top of a gold film. In this configuration, the vortices find it energetically easiest to settle into energy dimples in an arrangement like an egg crate—and make the material act as a Mott insulator, since the vortices won't move if the applied electric current is small.



experiment with and provides what looks like access to very complicated [many-body systems](#)," said Vinokur. "It looks a bit like magic."

As the name implies, many-body problems involve a large number of particles interacting; with current theory they are very difficult to model or understand.



When they applied a large enough electric current, however, the scientists saw a dynamic Mott transition as the system flipped to become a conducting metal; the properties of the material had changed as the current pushed it out of equilibrium.

The vortex system behaved exactly like an electronic Mott transition driven by temperature, said Valerii Vinokur, an Argonne Distinguished Fellow and corresponding author on the study. He and study co-author Tatyana Baturina, then at Argonne, analyzed the data and recognized the Mott behavior.

"This experimentally materializes the correspondence between quantum and classical physics," Vinokur said. "We can controllably induce a phase transition between a state of locked vortices to itinerant vortices by applying an electric current to the system," said Hans Hilgenkamp, head of the University of Twente research group. "Studying these [phase transitions](#) in our artificial systems is interesting in its own right, but may also provide further insight in the electronic transitions in real materials."

The system could further provide scientists with insight into two categories of physics that have been hard to understand: many-body systems and out-of-equilibrium systems.

"This is a classical system that which is easy to

"Furthermore, this system will be key to building a general understanding of out-of-equilibrium physics, which would be a major breakthrough in physics," Vinokur said.

The Department of Energy named five great basic energy scientific challenges of our time; one of them is understanding and controlling out-of-equilibrium phenomena. Equilibrium systems—where there's no energy moving around—are now understood quite well. But nearly everything in our lives involves energy flow, from photosynthesis to digestion to tropical cyclones, and we don't yet have the physics to describe it well. Scientists think a better understanding could lead to huge improvements in energy capture, batteries and energy storage, electronics and more.

As we seek to make electronics faster and smaller, Mott systems also offer a possible alternative to the silicon transistor. Since they can be flipped between conducting and insulating with small changes in voltage, they may be able to encode 1s

and 0s at smaller scales and higher accuracy than silicon transistors.

'Initially, we were studying the structures for completely different reasons, namely to investigate the effects of inhomogeneities on superconductivity," Hilgenkamp said. "After discussing with Valerii Vinokur at Argonne, we looked more specifically into our data and were quite amazed to see that it revealed so nicely the details of the transition between the state of locked and moving vortices. There are many ideas for follow up studies, and we look forward to our continued collaboration."

The results were printed in the study "Critical behavior at a dynamic vortex insulator-to-metal transition," released today in *Science*. Other co-authors are associated with the Siberian Branch of Russian Academy of Science, the Rome International Center for Materials Science Superstripes, Novosibirsk State University, the Moscow Institute of Physics and Technology and Queen Mary University of London.

More information: "Critical behavior at a dynamic vortex insulator-to-metal transition." *Science* 11 September 2015: [DOI: 10.1126/science.1260507](https://doi.org/10.1126/science.1260507)

Provided by University of Twente

APA citation: Team announces breakthrough observation of Mott transition in a superconductor (2015, September 11) retrieved 18 May 2021 from <https://phys.org/news/2015-09-team-breakthrough-mott-transition-superconductor.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.