

Scientists optimize breeding management for European minks

22 September 2015



European Mink. Credit: Tiit Maran

The European mink (*Mustela lutreola*) is one of the most endangered mammals in Europe. The reasons for its decline are the destruction of its habitat in riparian areas, competition with the alien American mink and historically, extensive hunting.

The European [mink](#) is often confused with the American mink (*Neovison vison*, previously *Mustela vison*), which has successfully established itself in Europe as an escapee from fur farms. The larger and more robust American mink has nearly completely replaced the European mink in its previous range.

Species protection projects all over Europe have so far faced the problem that European minks are difficult to breed in zoos. Captivity appears to have a negative effect on breeding success. But captive-bred individuals are needed in order to release and reintroduce the animal into protective zones. "The more we know about the physiology of European minks, the better we can respond to their needs," says lead author Franz Schwarzenberger from the Institute of Physiology, Pathophysiology and Experimental Endocrinology at the University of

Veterinary Medicine, Vienna.

Faecal samples yield reproductive data

Scientists from the Vetmeduni Vienna, in cooperation with the Endangered Species Research Lab of Tallinn Zoo, collected [faecal samples](#) from European mink and analysed them in Vienna. The [animals](#) are managed under the aegis of an EEP program (European Endangered Species Program). Female oestrus is usually determined by vaginal cytology. The aim of the study was to assess the validity of this method and to optimise diagnosis of ovulation and pregnancy.

"Using our non-invasive method, we were able to measure female oestrogen levels and generate a seasonal hormone profile. The results showed that oestrogen levels are higher at the time of ovulation. Such oestrogen peaks occur three to four times a year on average. The animals are polyoestrous. That means, during the breeding season they are fertile in regular intervals. In the past, females which had already been mated with no success were not mated again that same year. Our results reveal that mating can occur much more often," Schwarzenberger explains.

Looking for the perfect mate

European minks are solitary animals and extremely territorial in the wild, only approaching each other during the breeding season. In captivity, the animals are housed in large individual enclosures. "The exact time for mating is difficult to determine in a zoo because the animals attack each other if they aren't receptive. In order to increase the chances of fertilisation, the females are examined at regular intervals during the mating season. During mating, we also closely observe the behaviour of the animals, especially of the males," explains Astrid Nagl, first author of the study.

The Tallinn Zoo uses vaginal cytology to predict the

time of ovulation. This method does not always yield satisfactory results, however. "The data from the faecal analysis serve to augment the available information so that some females which had previously not been mated successfully also had offspring," Schwarzenberger reports.

Reintroduction in Europe not possible everywhere

The reintroduction of the European mink in Austria would not be easy. "In Austria, the American mink has replaced the European mink in aquatic and riparian zones," says Schwarzenberger. Releasing the European mink in this habitat would be tantamount to a death sentence, as the American minks would defend their territory and kill the European Mink. This makes reintroduction only possible in areas where no populations of American mink exist," says Schwarzenberger.

About 100–120 European mink live at Tallinn Zoo. The zoo's captive-bred animals are reintroduced to the wild on the Estonian islands of Hiiumaa and Saaremaa. Another promising reintroduction project can be found at Steinhuder Meer in northwest Germany.

More information: "Noninvasive monitoring of female reproductive hormone metabolites in the endangered European mink (*Mustela lutreola*)," *Theriogenology*, Available online 5 August 2015, ISSN 0093-691X, [dx.doi.org/10.1016/j.theriogenology.2015.07.023](https://doi.org/10.1016/j.theriogenology.2015.07.023)

Provided by University of Veterinary
Medicine—Vienna

APA citation: Scientists optimize breeding management for European minks (2015, September 22) retrieved 20 September 2020 from <https://phys.org/news/2015-09-scientists-optimize-european-minks.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.