

Arid forests provide refuge against cane toads and fires

28 September 2015, by Geoff Vivian, Sciencenetwork Wa



Northern spiny-tailed gecko. Credit: Ian Radford

Early research shows arid Kimberley woodlands may serve as a refuge for endemic fauna depleted by toxic cane toads (*Rhinella marina*) in other habitats.

Department of Parks and Wildlife fire ecologist Ian Radford led seven surveys of reptiles, frogs, invertebrates and mammals in the pindan woodland of Mirima National Park, near Kununurra, between 2008 and 2012.

They made four surveys before toads invaded the east Kimberley pindan, which is scrubland dominated by pindan wattle (*Acacia tumida*), and three after the incursion, noting population changes.

"The numbers of goannas did drop off, there's only about half to a third of the number of goannas after [cane toads](#) came through than there were beforehand," Dr Radford says.

"The numbers of snakes remained exactly the same."

He says they found an increase in some species

commonly eaten by goannas (*Varanus* spp.) and snakes, namely smaller skinks (*Carlia* spp.) up by 426 per cent and toadlets (*Uperoleia* spp.) up 306 per cent.

There was a 50 per cent decrease in [invertebrates](#), and a type of tiny sand-burrowing lizard (*Lerista griffin*) down by 87 per cent—which may have been due to more frogs and skinks eating them.

Dr Radford stressed the conclusions were tentative, as they had been unable to establish toad-free control areas throughout the survey period.

This would have established a clearer correlation between the influx of toads and changing numbers of other species, which may have also been influenced by other unknown environmental factors.



Trioda grassland. Credit: Hesperian

Notwithstanding this weakness in survey design, the team determined the pindan fauna numbers much less affected by the cane toad invasion than animals living near creeks and waterholes.

Dr Radford originally designed the survey to study differing fire regimes effects on the east Kimberley

pindan, as fires kill the dominant tree *A. tumida* which can then only re-establish itself from seed.

This involved studying the effects of hot wildfires, cooler prescribed burns, and no burning at all, on the region's flora.

He found the animals re-established themselves completely within one year of a hot wildfire, just as they do in the more arid savannah grasslands dominated by spinifex (*Triodia* spp).

However, the pindan wattle took four to five years to sprout from seed and reach its full height.

He found these destructive wildfires correlated with several years' build-up of unburned fuel—dead leaves, sticks and long grasses—over larger areas.

This article first appeared on [ScienceNetwork Western Australia](#) a science news website based at Scitech.

Provided by Science Network WA

APA citation: Arid forests provide refuge against cane toads and fires (2015, September 28) retrieved 20 May 2019 from <https://phys.org/news/2015-09-arid-forests-refuge-cane-toads.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.