

Global marine analysis suggests food chain collapse

12 October 2015



Credit: Tiago Fioreze / Wikipedia

A world-first global analysis of marine responses to climbing human CO₂ emissions has painted a grim picture of future fisheries and ocean ecosystems.

Published today in the journal *Proceedings of the National Academy of Sciences (PNAS)*, marine ecologists from the University of Adelaide say the expected ocean acidification and warming is likely to produce a reduction in diversity and numbers of various key [species](#) that underpin [marine ecosystems](#) around the world.

"This 'simplification' of our oceans will have profound consequences for our current way of life, particularly for coastal populations and those that rely on oceans for food and trade," says Associate Professor Ivan Nagelkerken, Australian Research Council (ARC) Future Fellow with the University's Environment Institute.

Associate Professor Nagelkerken and fellow University of Adelaide marine ecologist Professor Sean Connell have conducted a 'meta-analysis' of the data from 632 published experiments covering tropical to arctic waters, and a range of ecosystems

from coral reefs, through kelp forests to open oceans.

"We know relatively little about how climate change will affect the marine environment," says Professor Connell. "Until now, there has been almost total reliance on qualitative reviews and perspectives of potential global change. Where quantitative assessments exist, they typically focus on single stressors, single ecosystems or single species.

"This analysis combines the results of all these experiments to study the combined effects of multiple stressors on whole communities, including species interactions and different measures of responses to climate change."

The researchers found that there would be "limited scope" for acclimation to warmer waters and acidification. Very few species will escape the negative effects of increasing CO₂, with an expected large reduction in species diversity and abundance across the globe. One exception will be microorganisms, which are expected to increase in number and diversity.

From a total food web point of view, primary production from the smallest plankton is expected to increase in the warmer waters but this often doesn't translate into secondary production (the zooplankton and smaller fish) which shows decreased productivity under [ocean acidification](#).

"With higher metabolic rates in the warmer water, and therefore a greater demand for food, there is a mismatch with less food available for carnivores—the bigger fish that fisheries industries are based around," says Associate Professor Nagelkerken. "There will be a species collapse from the top of the food chain down."

The analysis also showed that with warmer waters or increased acidification or both, there would be deleterious impacts on habitat-forming species for

example coral, oysters and mussels. Any slight change in the health of habitats would have a broad impact on a wide range of species these reefs harbour.

Another finding was that acidification would lead to a decline in dimethylsulfide gas (DMS) production by ocean plankton which helps cloud formation and therefore in controlling the Earth's heat exchange.

More information: Global alteration of ocean ecosystem functioning due to increasing human CO₂ emissions, *PNAS*,
www.pnas.org/cgi/doi/10.1073/pnas.1510856112

Provided by University of Adelaide

APA citation: Global marine analysis suggests food chain collapse (2015, October 12) retrieved 15 June 2021 from <https://phys.org/news/2015-10-global-marine-analysis-food-chain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.