


Scientists track speed of powerful internal waves

13 October 2015



These two figures show the internal waves at Dongsha Island on April 23, 2010, as seen by the radar on TerraSAR-X in its conventional mode of operation (left) and in the experimental new mode that permits direct velocity measurements (right), with the measured surface velocities shown in color. Red and blue colors indicate surface velocities of about 0.5 m/s to the left and to the right, respectively. The shown area is 30 km × 80 km. Dongsha Island, which is about 2.7 km × 0.9 km (1.7 mi × 0.6 mi) in size, can be seen near the center of the image. Credit: German Aerospace Center (DLR) 2010.

For the first time researchers directly measured the speed of a wave located 80 meters below the ocean's surface from a single satellite image. The new technique developed by researchers from the University of Miami (UM) Rosenstiel School of

Marine and Atmospheric Science is a major advancement in the study of these skyscraper-high internal waves that rarely break the ocean surface.

"This is the first time internal wave velocities could be calculated from data acquired during a single overpass of a satellite," said Roland Romeiser, associate professor of ocean sciences at the UM Rosenstiel School. "This allows us to obtain more accurate information from a satellite that we could in the past."

Using a single satellite image collected at UM's Center for Southeastern Tropical Remote Sensing (CSTARS), the research team was able to determine that a roughly 60-meter high internal wave was traveling at a speed of three miles per hour (1.4 meters per second) near Dongsha Island in the South China Sea. The region is considered to have some of the most powerful internal waves on the planet.

"This is a significant breakthrough using a single image to determine the velocity of a wave below the surface," said Hans Graber, UM Rosenstiel School professor of ocean sciences and director of CSTARS. "This technology offers new opportunities to track the speed of ocean currents or objects moving on or below the [ocean surface](#)."

Radar satellites can detect the surface ripples produced by internal waves and the data collected allow researchers to calculate the speed of internal waves traveling below the surface. Prior to the development of this new technology, researchers would have to compare several images taken during multiple satellite overpasses to estimate internal wave velocities. The radar affixed to the German satellite TerraSAR-X is the first to measure velocities directly during a single overpass but with significant noise. Romeiser and Graber developed a new method to process the data that enhances the internal wave patterns to extract the velocities with unprecedented accuracy. CSTARS is the only

place besides the German Aerospace Center (DLR) that is capable of processing these types of images.

Internal waves move huge volumes of heat, salt, and nutrient rich-water across the ocean, which is important to fish, industrial fishing operations and the global climate. In addition, they are important to monitor for safe surface and sub-surface marine operations.

Graber was part of an international research team that spent seven years tracking the movements of internal waves to understand how these waves develop, move and dissipate underwater. The team discovered that [internal waves](#) are generated daily from internal tides, which also occur below the ocean surface, and grow larger as the water is pushed westward through the Luzon Strait into the South China Sea. Their findings were published in the May 7 issue of the journal *Nature*.

A research team led by Romeiser was the first to accurately measure currents from a space shuttle platform between islands off the Dutch coast and the first to make current measurements using the radar on the TerraSAR-X satellite.

The study, titled "Advanced Remote Sensing of Internal Waves by Spaceborne Along-Track InSAR—A Demonstration With TerraSAR-X," appears in the Dec. 2015 issue of the journal *Transactions on Geoscience and Remote Sensing*, a publication of the Institute of Electronic and Electrical Engineers (IEEE). The study's authors are Roland Romeiser and Hans Graber of the UM Rosenstiel School. The work was supported by grants from the U.S. Office of Naval Research.

More information: The study, titled "Advanced Remote Sensing of Internal Waves by Spaceborne Along-Track InSAR—A Demonstration With TerraSAR-X," appears in the Dec. 2015 issue of the journal *Transactions on Geoscience and Remote Sensing*, a publication of the Institute of Electronic and Electrical Engineers (IEEE).

Provided by University of Miami

APA citation: Scientists track speed of powerful internal waves (2015, October 13) retrieved 25 June 2019 from <https://phys.org/news/2015-10-scientists-track-powerful-internal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.