

Cyanobacteria can efficiently harvest sunlight with smart antenna

15 October 2015



Cyanobacteria growing in a multicultivator. Credit: Molecular Plant Biology Unit, University of Turku

Cyanobacteria have an interesting trick to harvest sunlight during light fluctuations. In darkness, the cells prepare for a subsequent increase in light intensity by adopting a larger light-harvesting antenna. Researchers of Wageningen University and the University of Turku, Finland describe the mechanism in a study that appeared online in Nature's *Scientific Reports*.

Cyanobacteria are ancient single-celled aquatic organisms, also known as blue-green algae: their blue color arises from their big light-harvesting pigment-protein complexes. These complexes contain blue pigments that are most sensitive to orange-green light, which can penetrate deeply into the water. Cyanobacteria contribute substantially to the global fixation of carbon dioxide, and knowledge about their photosynthetic strategies may contribute to original solutions for energy sustainability.


Regulatory mechanisms

The aquatic environment poses various challenges to the photosynthetic apparatus. In the upper water layers, one of the main challenges is the fluctuating incident light intensity. Waves and currents cause scattering of the light in waters and, in particular, suddenly increased [light intensities](#) can cause imbalances in the photosynthetic apparatus. It is well-known that the cells therefore have regulatory mechanisms to rapidly modulate

the amount of absorbed light by rearranging the light-harvesting machinery.

Natural light conditions

Due to the complexity of the photo-regulatory mechanisms, most of the research makes use of highly specific, well-defined artificial conditions, such as single-wavelength light and the presence of chemicals to regulate the energy flow. The new study uses 'natural' light conditions: a dark-to-light scheme typical of what the organisms experience when external factors move them between these regimes in the sea.



A 800ps fluorescence signature from cyanobacterial cells obtained with time-resolved fluorescence spectroscopy, the main technique used in this work.

Antenna

In darkness, the cells accumulate many electrons due to respiratory processes. These electrons can be re-used for the creation of energy-rich species during photosynthesis. The study shows that in

darkness, the cells seem to "get ready" for an increase in light intensity by adopting a larger light-harvesting antenna. In this way, during the first seconds after the [light](#) appears, they can immediately utilize the electron pool very efficiently to re-create energy-rich molecules.

More information: Volha Chukhutsina et al. Cyanobacterial Light-Harvesting Phycobilisomes Uncouple From Photosystem I During Dark-To-Light Transitions, *Scientific Reports* (2015). [DOI: 10.1038/srep14193](#)

Provided by Wageningen University
APA citation: Cyanobacteria can efficiently harvest sunlight with smart antenna (2015, October 15) retrieved 4 March 2021 from <https://phys.org/news/2015-10-cyanobacteria-efficiently-harvest-sunlight-smart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.