

# Extreme weather events in Chesapeake Bay give clues for future climate impacts

31 October 2015



The Chesapeake Bay area, with Baltimore to the north is shown. Image courtesy NASA Goddard Space Flight Center Landsat Data Continuity Mission Education and Public Outreach team (GSFC LDCM EPO). Credit: NASA Goddard Space Flight Center Landsat Data Continuity Mission Education and Public Outreach team (GSFC LDCM EPO).

For the millions of people who live in its expansive coastal areas, Chesapeake Bay provides an important source of income and recreational enjoyment. To protect the ecosystem and the livelihood of area residents, it is important to assess how climate variability and change will affect Chesapeake Bay's shallow water ecosystems and water quality.

The intensity, duration, and frequency of [extreme temperature](#)- and precipitation-based events are key components to understanding the climate of Chesapeake Bay.

Kari Pohl of the Center for Environmental Science at the University of Maryland and colleagues studied these components by calculating 26 extreme climate indices defined by the Expert Team on Climate Change Detection and Indices. She will report their findings at the Annual Meeting of the Geological Society of America on 1 November in Baltimore, Maryland, USA.

"A traditional view of historic climate change is to determine mean annual changes," Pohl explained. "However, organisms do not feel means; they feel the day-to-day variability and [extreme events](#), such as the frequency of warmer-than-normal days."

The goals of the project include reconstructing extreme climate changes from the recent past (1894-2014), using historically referenced data to assess near-future global climate model projections, and to ultimately use this analysis to investigate ecological problems in Chesapeake Bay, such as eelgrass diebacks.

The study saw changes that included an overall decrease in cold events, a higher probability to have a year without a cold spell, and an increase in the annual number of wet days. These extreme climate indices were strongly correlated to the shallow water environment, including streamflow and water temperature. These linkages will allow insights on how extreme changes could affect environmental boundaries and critical threshold events of vulnerable organisms.

Pohl hopes that studies such as this one "will enhance our general understanding of historical and future [extreme climate](#) variability, allowing policy-makers to make better-informed decisions for coastal communities."

**More information:** Paper No. 10: Chesapeake Bay Climate Extremes and Variability: a Recent Past, Present, and Near Future Analysis  
Abstract hyperlink:

[gsa.confex.com/gsa/2015AM/webp ...  
ram/Paper268364.html](https://gsa.confex.com/gsa/2015AM/webprogram/Paper268364.html)

Provided by Geological Society of America

APA citation: Extreme weather events in Chesapeake Bay give clues for future climate impacts (2015, October 31) retrieved 27 October 2021 from <https://phys.org/news/2015-10-extreme-weather-events-chesapeake-bay.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*