

High toxin level delays California crab season

5 November 2015



Fishermen stand on a boat loaded with crab pots, left, Thursday, Nov. 5, 2015, at Fisherman's Wharf in San Francisco. Wildlife authorities delayed the Dungeness crab season and closed the rock crab fishery for most of California on Thursday, just days after warning of dangerous levels of a neurotoxin linked to a massive algae bloom off the West Coast. (AP Photo/Ben Margot)

Wildlife authorities delayed the Dungeness crab season and closed the rock crab fishery for most of California on Thursday, just days after warning of dangerous levels of a neurotoxin linked to a massive algae bloom off the West Coast.

The state Fish and Game Commission voted 3-0 on the Dungeness delay and the rock crab closure north of the Santa Barbara-Ventura county line. The panel said crabbing would resume when toxin levels dropped but did not estimate when that might be.

The recreational Dungeness season was scheduled to start Saturday and the commercial season was set for Nov. 15. Rock [crabs](#) are caught year-round.

Officials on Tuesday warned people to avoid eating Dungeness and rock crabs. High levels of domoic acid have been found in crabs from the


Oregon border to the southern Santa Barbara County line, the Department of Public Health reported.

In severe poisoning cases, the neurotoxin can cause seizures, coma or death.

It was unclear how much impact the actions might have on California crabbing, which is estimated to bring in at least \$60 million commercially.

The toxin is linked to a vast algae bloom off the West Coast—which has seen unusually warm ocean temperatures as a result of El Nino, said Jordan Traverso, a spokeswoman at the California Department of Fish and Wildlife.

Such blooms are cyclical, but this summer surveyors aboard a National Oceanic and Atmospheric Administration research vessel said the algae bloom was one of the largest ever observed on the West Coast.



Crab pots wait to be loaded onto fishing boats Thursday, Nov. 5, 2015, at Fisherman's Wharf in San Francisco. Wildlife authorities delayed the Dungeness crab season and closed the rock crab fishery for most of California on Thursday, just days after warning of dangerous levels of a neurotoxin linked to a massive algae bloom off the West Coast. (AP Photo/Ben Margot)

The toxin has affected shellfish and sickened or killed seabirds, seals, dolphins and whales throughout the region.

Oregon state officials issued an advisory this week for all recreationally caught crab along the state's southern coast, from south of Coos Bay to California. Officials warned people to remove the viscera, or guts, before eating the crab meat.

Last month, Washington shellfish managers postponed the fall start of razor clam digging on ocean beaches, and all razor clamming remains closed along the Oregon coast because of high levels of domoic acid.



Crab pots wait to be loaded onto boats that are docked Thursday, Nov. 5, 2015, at Fisherman's Wharf in San Francisco. California officials are considering delaying the Dungeness crab season and closing the rock crab fishery just days after warning of dangerous levels of a neurotoxin linked to a massive algae bloom off the West Coast. (AP Photo/Ben Margot)

[Questions, answers about toxic West Coast crabs](#)

© 2015 The Associated Press. All rights reserved.

APA citation: High toxin level delays California crab season (2015, November 5) retrieved 2 December 2020 from <https://phys.org/news/2015-11-high-toxin-california-crab-season.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no

part may be reproduced without the written permission. The content is provided for information purposes only.