

Earth's climate more sensitive to CO2 than previously thought, study finds

16 November 2015



A modern trona from Lake Magadi, Kenya. Credit: David Tuttle

400 ppm. According to current projections, doubling the CO2 will result in a rise in the [global average temperature](#) of 3 degrees Centigrade. This new research suggests that the effects of CO2 on global warming may be underestimated.


"Take notice that carbon dioxide 50 million years ago may not have been as high as we once thought it was. We may reach that level in the next century, and so the [climate change](#) from that increase could be pretty severe, pretty dramatic. CO2 and other climate forcings may be more important for [global warming](#) than we realized."

Ancient climates on Earth may have been more sensitive to carbon dioxide than was previously thought, according to new research from Binghamton University.

A team of Binghamton University researchers including geology PhD student Elliot A. Jagniecki and professors Tim Lowenstein, David Jenkins and Robert Demicco examined nahcolite crystals found in Colorado's Green River Formation, formed 50 million years old during a hothouse [climate](#). They found that CO2 levels during this time may have been as low as 680 parts per million (ppm), nearly half the 1,125 ppm predicted by previous experiments. The new data suggests that past predictions significantly underestimate the impact of [greenhouse warming](#) and that Earth's climate may be more sensitive to increased carbon dioxide than was once thought, said Lowenstein.

"The significance of this is that CO2 50 million years ago may not have been as high as we once thought it was, but the climate back then was significantly warmer than it is today," said Lowenstein."

CO2 levels in the atmosphere today have reached



Lowenstein wants to look at nahcolite deposits in China to confirm the results found in Colorado.

More information: Elliot A. Jagniecki et al. Eocene atmospheric CO from the nahcolite proxy , *Geology* (2015). [DOI: 10.1130/G36886.1](https://doi.org/10.1130/G36886.1)

Provided by Binghamton University

A nahcolite from the Eocene Green River Formation.
Credit: Timothy Lowenstein

The only direct measurement of carbon dioxide is from ice cores, which only go back less than 1 million years. Lowenstein and his team are trying to develop ways to estimate ancient [carbon dioxide](#) in the atmosphere using indirect proxies. He said that their approach is different than any ever undertaken.

"These are direct chemical measurements that are based on equilibrium thermodynamics," he said. "These are direct laboratory experiments, so I think they're really reliable."

APA citation: Earth's climate more sensitive to CO₂ than previously thought, study finds (2015, November 16) retrieved 24 June 2021 from <https://phys.org/news/2015-11-earth-climate-sensitive-co2-previously.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.