

# Dating historic activity at Oso site shows recurring major landslides

23 December 2015

The large, fast-moving mudslide that buried much of Oso, Washington in March 2014 was the deadliest landslide in U.S. history. Since then, it's been revealed that this area has experienced major slides before, but it's not known how long ago they occurred.

University of Washington geologists analyzed woody debris buried in earlier slides and used radiocarbon dating to map the history of activity at the site. The findings, published online in the journal *Geology*, show that a massive nearby slide happened around 500 years ago, and not thousands of years ago as some had believed.

"The soil in this area is all glacial material, so one hypothesis is the material could have fallen apart in a series of large landslides soon after the ice retreated, thousands of years ago," said corresponding author Sean LaHusen, a UW doctoral student in Earth and space sciences. "We found that that's not the case—in fact, landslides have been continuing in recent history."

The study establishes a new method to date all the previous landslides at a particular location. The method shows that the slopes in the area around Oso have collapsed on average once every 500 years, and at a higher rate of about once every 140 years over the past 2,000 years.

"This was well known as an area of hillslope instability, but the question was: 'Were the larger slides thousands of years old or hundreds of years old?' Now we can say that many of them are hundreds of years old," said co-author Alison Duvall, a UW assistant professor of Earth and space sciences.

LaHusen had not yet begun his graduate studies when he asked about studying the history of geologic activity at the Oso site. In late summer of 2014, the researchers began their work wading along riverbanks to look for preserved branches or

trees that could be used to date previous landslides.

"When you have a large, catastrophic landslide, it can often uproot living trees which kills them and also encapsulates them in the landslide mass," Duvall said. "If you can find them in the landslide mass, you can assume that they were killed by the landslide, and thus you can date when the landslide occurred."

The team managed to unearth samples of wood buried in the Rowan landslide, just downstream of the Oso site, and the Headache Creek landslide, just upriver of the 2014 slide. Results from several debris samples show that the Rowan landslide, approximately five times the size of the Oso slide, took place just 300 to 694 years ago. The Headache Creek landslide is within a couple hundred years of 6,000 years old.

Previous UW research had shown a history of geologic activity at the Oso site, including previous major landslides and a recent small slide at the same slope that collapsed in 2014. But while the position of past slides and degree of surface erosion can show the order that the older slides happened, it has not been possible to give a date for the past events.

The new study uses the radiocarbon dates for two slides to establish a roughness curve to date other events along a 3.7-mile (6-kilometer) stretch of the north fork of the Stillaguamish River. A roughness curve uses the amount of surface erosion to establish each slide's age. The two dates put firm limits on the curve, so that other nearby slides can be dated from their roughness characteristics without having to find material buried inside each mass of soil.

"This is the first time this calibrated surface dating method has been used for landslide chronologies, and it seems to work really well," LaHusen said. "It can provide some information about how often

these events recur, which is the first step toward a regional risk analysis."

Applying the new method for other locations would require gathering samples for each area, they cautioned, because each site has its own soil composition and erosion characteristics.

It's not known whether the findings for the Oso site's history would apply to other parts of the Stillaguamish River, Duvall said, or to other places in Washington state. The researchers are still studying debris from other locations. But the results do have implications for the immediate area.

"It suggests that the Oso [landslide](#) was not so much of an anomaly," Duvall said.

She and LaHusen are also working with the UW's M-9 Project, which is studying hazards from magnitude-9 earthquakes along the Cascadia subduction zone. They would like to learn whether landslides across Washington state coincided with past earthquakes, and use simulations of future shaking to predict which places in the state are most vulnerable to earthquake-triggered landslides.

**More information:** Sean R. LaHusen et al.

Surface roughness dating of long-runout landslides near Oso, Washington (USA), reveals persistent postglacial hillslope instability, *Geology* (2015).

[DOI: 10.1130/G37267.1](https://doi.org/10.1130/G37267.1)

Provided by University of Washington

APA citation: Dating historic activity at Oso site shows recurring major landslides (2015, December 23) retrieved 27 February 2021 from <https://phys.org/news/2015-12-dating-historic-oso-site-recurring.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*