

Graphene becomes superconductive—Electrons with 'no mass' flow with 'no resistance'

16 February 2016




Fig1. Crystal structure of Ca-intercalated bilayer graphene fabricated on SiC substrate. Insertion of Ca atoms between two graphene layers causes the superconductivity.

Graphene is a single-atomic carbon sheet with a hexagonal honeycomb network (Fig. 1). Electrons in graphene take a special electronic state called Dirac-cone where they behave as if they have no mass. This allows them to flow at very high speed, giving graphene a very high level of electrical conductivity.

This is significant because electrons with no mass flowing with no resistance in graphene could lead to the realization of an ultimately high-speed nano electronic device.

The collaborative team of Tohoku University and the University of Tokyo has developed a method to grow [high-quality graphene](#) on a silicon carbide (SiC) crystal by controlling the number of [graphene sheets](#). The team fabricated bilayer graphene with this method and then inserted calcium (Ca) atoms between the two graphene layers like a sandwich (Fig. 1).

They measured the [electrical conductivity](#) with the micro four-point probe method and found that the electrical resistivity rapidly drops at around 4 K (-269 °C), indicative of an emergence of superconductivity (Fig. 2).

The team also found that neither genuine bilayer graphene nor lithium-intercalated bilayer graphene shows superconductivity, indicating that the superconductivity is driven by the electron transfer from Ca atoms to graphene sheets.

The success in fabricating superconducting graphene is expected to greatly impact both the basic and applied researches of graphene.




Fig.2 Temperature dependence of electrical resistivity of Ca-intercalated bilayer graphene, measured by the micro four-point-probe method (inset). The resistivity shows a rapid decrease at around 4 K and reaches "zero" at 2 K, showing the emergence of superconductivity.

It is currently not clear what phenomenon takes place when the Dirac electrons with no mass become superconductive with no resistance. But based on the latest study results, further experimental and theoretical investigations would help to unravel the properties of superconducting graphene.

The superconducting transition temperature (T_c) observed in this study on Ca-intercalated [bilayer graphene](#) is still low (4 K). This prompts further studies into ways to increase T_c , for example, by replacing Ca with other metals and alloys, or changing the number of graphene sheets.

From the application point of view, the latest results pave the way for the further development of ultrahigh-speed superconducting nano devices such as a quantum computing device, which utilizes superconducting graphene in its integrated circuit.

More information: Satoru Ichinokura et al. Superconducting Calcium-Intercalated Bilayer Graphene, *ACS Nano* (2016). [DOI: 10.1021/acsnano.5b07848](#)

Provided by Tohoku University

APA citation: Graphene becomes superconductive—Electrons with 'no mass' flow with 'no resistance' (2016, February 16) retrieved 16 January 2021 from <https://phys.org/news/2016-02-graphene-superconductiveelectrons-mass-resistance.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.