

Fungi are at the root of tropical forest diversity—or lack thereof, study finds

February 22 2016, by Diana Yates



The study focused on tropical forest patches dominated by one tree species, *Oreomunnea mexicana*, pictured. Credit: James Dalling

The types of beneficial fungi that associate with tree roots can alter the fate of a patch of tropical forest, boosting plant diversity or, conversely,

giving one tree species a distinct advantage over many others, researchers report.

Their study, reported in the journal *Ecology Letters*, sought to explain a baffling phenomenon in some tropical forests: Small patches of "monodominant forest," where one species makes up more than 60 percent of the [trees](#), form islands of low diversity in the otherwise highly diverse tropical forest growing all around them.

The new study focused on mountain forests in Panama that harbor hundreds of tree species, but which include small patches dominated by the tree species *Oreomunnea mexicana*.

"Tropical ecologists are puzzled by how so many species co-occur in a tropical forest," said University of Illinois plant biology professor James Dalling, who led the study with graduate student Adriana Corrales and collaborators from Washington University in St. Louis and the Smithsonian Tropical Research Institute in Panama. "If one tree species is a slightly better competitor in a particular environment, you would expect its population to increase and gradually exclude other species."

That doesn't happen often in tropical forests, however, he said. Diversity remains high, and patches dominated by a single species are rare. Understanding how monodominant forests arise and persist could help explain how tropical forests otherwise maintain their remarkable diversity, he said.

The researchers focused on two types of fungi that form symbiotic relationships with trees: arbuscular mycorrhizas and ectomycorrhizas. Arbuscular mycorrhizas grow inside the roots of many different tree species, supplying phosphorous to their tree hosts. Ectomycorrhizas grow on the surface of tree roots and draw [nitrogen](#) from the soil, some of which they exchange for sugars from the trees. Ectomycorrhizas

cooperate with only a few tree species - 6 percent or less of those that grow in tropical forests.

Previous studies found that arbuscular mycorrhizas commonly occur in the most diverse tropical forests, while ectomycorrhizal fungi dominate low-diversity patches.

"When you walk in a patch of forest where 70 percent of the trees belong to a single species that also happens to be an ectomycorrhizal-associated tree, it makes you think there is something going on with the fungi that could be mediating the formation of these monodominant forests," Corrales said.

The researchers tested three hypotheses to explain the high abundance of *Oreomunnea*. First, they tested the idea that *Oreomunnea* trees are better able to resist species-specific pathogens than trees growing in more diverse forest areas.

"We were expecting that *Oreomunnea* seedlings would grow better in soil coming from beneath other *Oreomunnea* trees, because that's how the tree grows in nature," Corrales said. "But we found the opposite: The *Oreomunnea* suffered more from pathogen infection when grown in soil from the same species than in soil from other species."

The researchers next tested whether mature *Oreomunnea* trees supported nearby *Oreomunnea* seedlings by sending sugars to them via a shared network of ectomycorrhizal fungi. But they found no evidence of cooperation between the trees.

"The seedlings that were isolated from the ectomycorrhizas of other *Oreomunnea* trees grew better than those that were in contact with the fungi from other trees of the same species," Corrales said.

In a third set of experiments, the team looked at the availability of nitrogen inside and outside the *Oreomunnea* patches.

"We saw that inorganic nitrogen was much higher outside than inside the patches," Corrales said. Tree species that normally grow outside the patches did well on the high-nitrogen soils, but suffered when transplanted inside the *Oreomunnea* patches. A look at the nitrogen isotopes in the fungi, soils and in the seedlings' leaves revealed the underlying mechanism by which the fungi influenced the species growing inside and outside the *Oreomunnea* patches.

The team found evidence consistent with ectomycorrhizal uptake of nitrogen directly from decomposing material in the soil. These fungi make some of their nitrogen available to the *Oreomunnea* trees while starving other plants and soil microbes of this essential nutrient, Corrales said. The lack of adequate nitrogen means bacteria and fungi are unable to break down organic matter in the soil, causing most other trees to suffer because they depend on the nitrogen supplied by microbial decomposers, she said.

"We found a novel mechanism that can explain why certain [tree species](#) in [tropical forests](#) are highly abundant, and that is because their [fungi](#) provide them with a source of nitrogen that is not accessible to competing species," Dalling said. "So they have an advantage because their competitors are now starved of nitrogen."

Researchers have found recently that similar processes can occur in temperate forests, but this is the first study to link this process to tropical forest monodominance, Dalling said.

More information: Adriana Corrales et al. An ectomycorrhizal nitrogen economy facilitates monodominance in a neotropical forest, *Ecology Letters* (2016). [DOI: 10.1111/ele.12570](https://doi.org/10.1111/ele.12570)

Provided by University of Illinois at Urbana-Champaign

Citation: Fungi are at the root of tropical forest diversity—or lack thereof, study finds (2016, February 22) retrieved 26 April 2024 from <https://phys.org/news/2016-02-fungi-root-tropical-forest-diversityor.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.