


Heat and light get larger at the nanoscale

31 March 2016




A video of the high-precision micro-electromechanical system (MEMS) used to control the distance between two beams at different temperatures. The video is taken under a high magnification microscope. The whole video frame dimension is comparable to the diameter of a strand of human hair. Credit: Raphael St-Gelais, Lipson Nanophotonics Group, Columbia Engineering

In a new study recently published in *Nature Nanotechnology*, researchers from Columbia Engineering, Cornell, and Stanford have demonstrated heat transfer can be made 100 times stronger than has been predicted, simply by bringing two objects extremely close—at nanoscale distances—without touching. Led by Columbia Engineering's Michal Lipson and Stanford Engineering's Shanhui Fan, the team used custom-made ultra-high precision micro-mechanical displacement controllers to achieve heat transfer using light at the largest magnitude reported to date between two parallel objects.

"At separations as small as 40 nanometers, we achieved almost a 100-fold enhancement of heat transfer compared to classical predictions," says Lipson, Eugene Higgins Professor of Electrical

Engineering and professor of applied physics. "This is very exciting as it means that light could now become a dominant heat transfer channel between objects that usually exchange heat mostly through conduction or convection. And, while other teams have demonstrated heat transfer using light at the nanoscale before, we are the first to reach performances that could be used for energy applications, such as directly converting heat to electricity using [photovoltaic cells](#)."

All objects in our environment exchange heat with their surroundings using light. This includes the light coming at us from the sun, the glowing red color of the heating element inside our toaster ovens, or the "night vision" cameras that enable image recording even in complete darkness. But heat exchange using light is usually very weak compared to what can be achieved by conduction (i.e., by simply putting two objects in contact with each other) or by convection (i.e., using hot air). Radiative heat transfer at nanoscale distances, while theorized, has been especially challenging to achieve because of the difficulty of maintaining large thermal gradients over nanometer-scale distances while avoiding other heat transfer mechanisms like conduction.



A schematic of two beams at different temperatures exchanging heat using light. In the situation when the beams are far from each other (left), heat transfer resulting from thermal radiation is small. When the beams are brought very close from each other (right) heat transfer becomes almost 100 times larger than predicted by conventional thermal radiation laws. Credit: Raphael St-Gelais, Lipson Nanophotonics Group, Columbia Engineering

Lipson's team was able to bring objects at different temperatures very close to each other—at distances smaller than 100 nanometers, or 1/1000th of the diameter of a strand of human hair. They were able to demonstrate near-field radiative heat transfer between parallel SiC (silicon carbide) nanobeams in the deep sub-wavelength regime. They used a high-precision micro-electromechanical system (MEMS) to control the distance between the beams and exploited the mechanical stability of nanobeams under high tensile stress to minimize thermal buckling effects, thus keeping control of the nanometer-scale separation even at large thermal gradients.

Using this approach, the team was able to bring two parallel objects at different temperatures to distances as small as 42 nm without touching. In this case they observed that the heat transfer between the objects was close to 100 times stronger than what is predicted by conventional thermal radiation laws (i.e. "blackbody radiation"). They were able to repeat this experiment for temperature differences as high as 260°C (500°F) between the two objects. Such high temperature difference is especially important for energy conversion applications since, in these cases, the conversion efficiency is always proportional to the thermal difference between the hot and the cold objects involved.

"An important implication of our work is that thermal radiation can now be used as a dominant heat transfer mechanism between objects at different temperatures," explains Raphael St-Gelais, the study's lead author and postdoctoral fellow working with Lipson at Columbia Engineering. "This means that we can control heat flow with a lot of the same techniques we have for manipulating light. This is a big deal since there are a lot of interesting things we can do with light, such as converting it to electricity using photovoltaic cells."

St-Gelais and Linxiao Zhu, who co-authored the study and is a PhD candidate in Fan's group at Stanford, note that the team's approach can be scaled up to a larger effective area by simply arraying several nanobeams—on top of a

photovoltaic cell, for example—and by individually controlling their out-of-plane displacement using MEMS actuators. The researchers are now looking at applying their same approach for ultra-high-precision displacement control, this time with an actual photovoltaic cell to generate electricity directly from heat.

"This very strong, non-contact, [heat transfer](#) channel could be used for controlling the temperature of delicate nano devices that cannot be touched, or for very efficiently converting heat to electricity by radiating large amounts of heat from a hot [object](#) to a photovoltaic cell in its extreme proximity," Lipson adds. "And if we can shine a large amount of heat in the form of light from a hot object to a photovoltaic cell, we could potentially create compact modules for direct conversion of heat to electrical power. These modules could be used inside cars, for instance, to convert wasted heat from the combustion engine back to useful electrical power. We could also use them in our homes to generate electricity from alternative energy sources such as biofuels and stored solar energy."

More information: Raphael St-Gelais et al. Near-field radiative heat transfer between parallel structures in the deep subwavelength regime, *Nature Nanotechnology* (2016). [DOI: 10.1038/nnano.2016.20](#)

Provided by Columbia University School of Engineering and Applied Science

APA citation: Heat and light get larger at the nanoscale (2016, March 31) retrieved 15 June 2019 from <https://phys.org/news/2016-03-larger-nanoscale.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.