


# Venus Express' swansong experiment sheds light on Venus' polar atmosphere

April 19 2016

---



Visualisation of Venus Express during the aerobraking manoeuvre, during which the spacecraft orbited Venus at an altitude of around 130 km from 18 June to 11 July 2014. In the month before, the altitude was gradually reduced from around 200 km to 130 km. Credit: ESA - C. Carreau


Some of the final results sent back by ESA's Venus Express before it plummeted down through the planet's atmosphere have revealed it to be rippling with atmospheric waves – and, at an average temperature of

-157°C, colder than anywhere on Earth.

As well as telling us much about Venus' previously-unexplored polar regions and improving our knowledge of our planetary neighbour, the experiment holds great promise for ESA's ExoMars mission, which is currently winging its way to the Red Planet. The findings were published in the journal *Nature Physics* on 11 April 2016.

ESA's Venus Express arrived at Venus in 2006. It spent eight years exploring the planet from orbit, vastly outliving the mission's planned duration of 500 days, before running out of fuel. The probe then began its descent, dipping further and further into Venus' [atmosphere](#), before the mission lost contact with Earth (November 2014) and officially ended (December 2014).

However, Venus Express was industrious to the end; low altitude orbits were carried out during the final months of the mission, taking the spacecraft deep enough to experience measurable drag from the atmosphere. Using its onboard accelerometers, the spacecraft measured the deceleration it experienced as it pushed through the planet's upper atmosphere – something known as aerobraking.



This figure shows the density of Venus' atmosphere in the northern polar regions at altitudes of 130 to 190 km. All data points were gathered during different phases of the Venus Express Atmospheric Drag Experiment (VExADE), performed between 2008-2013 (values above 165 km) and from 24 June to 11 July 2014 (values below 140 km); the black dots to the lower right were from the aerobraking phase (AER), the black dots to the upper left from the Precise Orbit Determination phase (POD), and the grey dots from torque measurements (TRQ). Each coloured line represents a different scientific model of Venus' atmosphere. The dark blue line shows a model based on data from NASA's Pioneer Venus spacecraft, dubbed VTS3 (Hedin et al., 1983), which uses observations of Venus' equatorial latitudes gathered from 1978-1980 (extrapolated to the poles). The cyan line corresponds to another reference model of Venus' neutral upper atmosphere based on Pioneer Venus, named Venus International Reference Atmosphere (VIRA, Keating et al., 1985). The red

line corresponds to a model (Venus Polar Atmosphere Model) currently being developed by Ingo Müller-Wodarg. This model seeks to bridge the data gap shown in the figure from 140-165 km and present a unified vertical density profile for Venus' upper polar atmosphere. Credit: Figure courtesy of I. Müller-Wodarg (Imperial College London, UK)

"Aerobraking uses [atmospheric drag](#) to slow down a spacecraft, so we were able to use the accelerometer measurements to explore the density of Venus' atmosphere," said Ingo Müller-Wodarg of Imperial College London, UK, lead author of the study. "None of Venus Express' instruments were actually designed to make such in-situ atmosphere observations. We only realised in 2006 – after launch! – that we could use the Venus Express spacecraft as a whole to do more science."

When Müller-Wodarg and colleagues gathered their observations Venus Express was orbiting at an altitude of between 130 and 140 kilometres near Venus' polar regions, in a portion of Venus' atmosphere that had never before been studied in situ.


Previously, our understanding of Venus' polar atmosphere was based on observations gathered by NASA's Pioneer Venus probe in the late 1970s. These were of other parts of Venus' atmosphere, near the equator, but extrapolated to the poles to form a complete atmospheric reference model.

These new measurements, taken as part of the Venus Express Atmospheric Drag Experiment (VExADE) from 24 June to 11 July 2014, have now directly tested this model – and reveal several surprises.

For one, the polar atmosphere is up to 70 degrees colder than expected, with an [average temperature](#) of  $-157^{\circ}\text{C}$  (114 K). Recent temperature

measurements by Venus Express' SPICAV instrument (SPectroscopy for the Investigation of the Characteristics of the Atmosphere of Venus) are in agreement with this finding.

The polar atmosphere is also not as dense as expected; at 130 and 140 km in altitude, it is 22% and 40% less dense than predicted, respectively. When extrapolated upward in the atmosphere, these differences are consistent with those measured previously by VExADE at 180 km, where densities were found to be lower by almost a factor of two.



This frame shows a visualization of raw data from the Venus Express Atmospheric Drag Experiment (VExADE), performed from 24 June to 11 July 2014, at altitudes of 130-140 km in the atmosphere of Venus. The black lines

show 16 of the spacecraft's 18 orbital trajectories from that period. The grey background is a normalised map of the atmospheric gravity waves that were detected. The non-uniformity represents density perturbations in Venus' polar atmosphere; darker patches are less dense, and lighter patches more dense, than their surroundings. The average density perturbation amplitude is around 10% of the mean background density. The results of the VExADE experiment, reported in *Nature Physics* (Müller-Wodarg et al., 2016), showed that strong atmospheric gravity waves dominate the polar regions of Venus' atmosphere. Credit: ESA/Venus Express/VExADE/Müller-Wodarg et al., 2016

"This is in-line with our temperature findings, and shows that the existing model paints an overly simplistic picture of Venus' [upper atmosphere](#)," added Müller-Wodarg. "These lower densities could be at least partly due to Venus' polar vortices, which are strong wind systems sitting near the planet's poles. Atmospheric winds may be making the density structure both more complicated and more interesting!"

Additionally, the polar region was found to be dominated by strong atmospheric waves, a phenomenon thought to be key in shaping planetary atmospheres – including our own.

"By studying how the atmospheric densities changed and were perturbed over time, we found two different types of wave: Atmospheric [gravity waves](#) and planetary waves," explained co-author Sean Bruinsma of the Centre National D'Etudes Spatiales (CNES), France. "These waves are tricky to study, as you need to be within the atmosphere of the planet itself to measure them properly. Observations from afar can only tell us so much."

Atmospheric gravity waves are similar to waves we see in the ocean, or when throwing stones in a pond, only they travel vertically rather than horizontally. They are essentially a ripple in the density of a planetary

atmosphere – they travel from lower to higher altitudes and, as density decreases with altitude, become stronger as they rise. The second type, planetary waves, are associated with a planet's spin as it turns on its axis; these are larger-scale waves with periods of several days.

We experience both types on Earth. Atmospheric gravity waves interfere with weather and cause turbulence, while planetary waves can affect entire weather and pressure systems. Both are known to transfer energy and momentum from one region to another, and so are likely to be hugely influential in shaping the characteristics of a planetary atmosphere.

"We found atmospheric gravity waves to be dominant in Venus' polar atmosphere," added Bruinsma. "Venus Express experienced them as a kind of turbulence, a bit like the vibrations you feel when an aeroplane flies through a rough patch. If we flew through Venus' atmosphere at those heights we wouldn't feel them because the atmosphere just isn't dense enough, but Venus Express' instruments were sensitive enough to detect them."

Venus Express found atmospheric waves at an altitude of 130-140 km that the team think originated from the upper cloud layer in Venus' atmosphere, which sits at and below altitudes of approximately 90 km, and a planetary wave that oscillated with a period of five days. "We checked carefully to ensure that the [waves](#) weren't an artefact of our processing," said co-author Jean-Charles Marty, also of CNES.

This is not just a first for Venus Express; while the aerobraking technique has been used for Earth satellites, and was previously used on NASA-led missions to Mars and Venus, it had never before been used on any ESA planetary mission.

However, ESA's ExoMars Trace Gas Orbiter, which launched earlier this

year, will use a similar technique. "During this activity we will extract similar data about Mars' atmosphere as we did at Venus," added Håkan Svedhem, project scientist for ESA's ExoMars 2016 and Venus Express missions.

"For Mars, the aerobraking phase would last longer than on Venus, for about a year, so we'd get a full dataset of Mars' atmospheric densities and how they vary with season and distance from the Sun," added Svedhem. "This information isn't just relevant to scientists; it's crucial for engineering purposes as well. The Venus study was a highly successful test of a technique that could now be applied to Mars on a larger scale – and to future missions after that."

**More information:** Ingo C. F. Müller-Wodarg et al. In situ observations of waves in Venus's polar lower thermosphere with Venus Express aerobraking, *Nature Physics* (2016). [DOI: 10.1038/nphys3733](https://doi.org/10.1038/nphys3733)

Provided by European Space Agency

Citation: Venus Express' swansong experiment sheds light on Venus' polar atmosphere (2016, April 19) retrieved 26 April 2024 from <https://phys.org/news/2016-04-venus-swansong-polar-atmosphere.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.