

Insulating layer of air above the Greenland ice sheet reduces precipitation

April 29 2016



Max Berkelhammer digs a snow pit near Summit Station Greenland to collect samples for isotopic analysis at sled-towed research lab that breaks up the otherwise featureless snowy landscape. Credit: David Noone, Oregon State University

The Earth's climate has been warming, but even though the Greenland

ice sheet is melting rapidly in the coastal regions, there are large parts of the ice sheet (40 percent) where there has hardly been any melting on the surface. A warmer climate usually also means that there is more precipitation, but there has been no increase in the amount of precipitation on the ice sheet. New research with participation of the Niels Bohr Institute shows that this is due to an insulating layer of air that forms near the surface during the winter, which insulates the ice sheet from the upper atmosphere and reduces both evaporation and precipitation. The results are published in the scientific journal, *Science Advances*.

When the [air](#) gets warmer, it can hold more [water](#) vapour. The laws of physics state that for every degree the temperature rises in the air, the water content can increase by 7 percent, so the air in tropical regions is much more humid than in Arctic Regions.

The amount of precipitation is therefore linked to temperature. But researchers with participation of the Niels Bohr Institute discovered that even though the temperature has risen over the last 10 years, the amount of precipitation over the central parts of the Greenland [ice sheet](#) has not increased. Measurements showed that the amount of precipitation actually decreased. Why so?

Because of low-lying clouds and driving snow, direct observations of the ice sheet surface can be very uncertain. Instead, an international team of researchers from the United States and the Niels Bohr Institute in Denmark have studied the [atmosphere](#) over the 3½ km thick ice sheet.

Finding answers in the air

"We decided to investigate whether you could find the answer in the atmosphere above the ice sheet by measuring the atmospheric processes directly. We therefore took measurements of the water vapour in the

atmosphere for three years," explains Hans Christian Steen-Larsen, a postdoc at the Centre for Ice and Climate at the Niels Bohr Institute at the University of Copenhagen.




Fig. 2. Vertical profiles of the isotopic ratio ($\delta^{18}\text{O}$) and humidity at Summit Camp. (Top) Isotopic ratio profile. (Bottom) Water vapor mixing ratio profile. The seasonal profiles were generated as the average of all hourly profiles from the respective time window and reported relative to the value at the top inlet. Red (green) lines are generated from profiles during stable (unstable) periods (fig. S3), and the error envelope represents the 66th and 90th percentiles. The dotted line is the average of all profiles. Blue dots below the profiles are the average isotopic ratios measured from the firn interstitial vapor. The error envelop for the firn vapor is 1 SD.

MS no: RA1501704/LB	Emp name: JAfan	Date / Time: 4-12-2016 / 5:2	PE's:	AA's:	Comments: 3	Art no: Fig. 2
------------------------	--------------------	---------------------------------	-------	-------	----------------	--------------------------

Teaser: The surface of the Greenland ice sheet becomes isolated from the atmosphere during the winter, which prohibits the exchange (loss or gain) of ice mass.

The curve at the isotopic composition of the water vapour between the snow surface and a height of 40 m shows that at a height of 2-4 meters there is a minimum of ^{18}O . So there is a boundary layer that separates the lower layer and the upper layer in the atmosphere. Credit: Berkelhammer et al.

Water consists of molecules with one oxygen atom and two hydrogen atoms. But the atoms can vary a bit and have more or fewer particles in the nucleus, which are called isotopes. The researchers measured the stable water isotopes in the water vapour in the atmosphere up to a height of 40 meters above the surface of the ice sheet. They analysed the water vapour and measured the ratio between the most common form of oxygen, ^{16}O and the slightly heavier oxygen, ^{18}O . This isotope ratio says something about where the water vapour came from and they got some very surprising results.

It turns out that there are very stable weather systems in the atmosphere over the ice sheet during the winter. Close to the surface it is very cold and dry and the air is very stagnant. About 100 meters up, the air is more dynamic and warmer. But it is not a smooth transition.

Discovering a new phenomena

"By measuring the isotopic composition of the water vapour, we discovered that there was a very clear distinction between the water vapour that was associated with the ice and the water vapour that was blown in higher up in the atmosphere. There was a high content of the oxygen isotope ^{18}O both in the lowest [layer](#) of air near to the ice and in the higher layer of air. But in between the two air layers at a height of 2-4 meters there was a layer with a lower content of ^{18}O . So there was a

boundary layer that separated the two layers in the atmosphere. We were thus able to see a direct disconnection between the surface of the ice and the atmosphere above it - it would not have been possible to discover this if we had only looked at the water vapour amount in the air," explains Hans Christian Steen-Larsen.



Hans Christian Steen-Larsen is wearing climbing equipment in minus 40 degrees at Summit on the ice sheet. Credit: David Schneider, NCAR

This disconnect between the two layers explains why there is less precipitation than expected. Above the boundary layer, the air contains water and can form a fog, but because the fog is simply too thin and the particles are too small, the fog does not get dense enough to fall as precipitation, so it simply evaporates again. At the same time, the thin layer of fog puts a layer on the surface, preventing evaporation from the surface of the ice.

"We have simply discovered an important process that helps to explain why there is no relationship between temperature and precipitation on the Greenland ice sheet," says Hans Christian Steen-Larsen.

Better climate models

The results were further tested by comparing them with data from ice cores. The Greenland ice sheet was formed by snow that falls year after year and remains and is gradually compressed into ice. By drilling ice cores through the kilometers thick ice sheet, researchers can clearly see each annual layer and they can measure what the temperature was in the precipitation that originally fell as snow. In this way, they can discover what the climate was like over more than a hundred thousand years - how warm it was and how much precipitation fell.

"If we look at the data from the last 11,000 years, where we have had a warm interglacial period, we can see that the temperature and the amount of precipitation have not always followed each other. This missing link has been difficult to explain, but we can now do this with our discovery of this phenomenon. This means that you can use data

from the ice cores that tells us about past climate variations to improve the [climate models](#) for predicting future climate chance," concludes Hans Christian Steen-Larsen.

More information: Surface-atmosphere decoupling limits accumulation at Summit, Greenland, *Science Advances*, DOI: 10.1126/sciadv.1501704,
advances.sciencemag.org/content/2/4/e1501704

Provided by Niels Bohr Institute

Citation: Insulating layer of air above the Greenland ice sheet reduces precipitation (2016, April 29) retrieved 20 September 2024 from <https://phys.org/news/2016-04-insulating-layer-air-greenland-ice.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.