


Scientists observe supermassive black hole feeding on cold gas

8 June 2016



The cosmic weather report, as illustrated in this artist concept, calls for condensing clouds of cold molecular gas around the Abell 2597 Brightest Cluster Galaxy. The clouds condense out of the hot, ionized gas that suffuses the space between the galaxies in this cluster. New ALMA data show that these clouds are raining in on the galaxy, plunging toward the supermassive black hole at its center. Credit: NRAO/AUI/NSF; D. Berry / SkyWorks; ALMA (ESO/NAOJ/NRAO)

At the center of a galaxy cluster, 1 billion light years from Earth, a voracious, supermassive black hole is preparing for a chilly feast.

For the first time, astronomers have detected

billowy clouds of cold, clumpy gas streaming toward a black hole, at the center of a massive galaxy cluster. The clouds are traveling at speeds of up to 355 kilometers per second—that's almost 800,000 miles per hour—and may be only 150 light years away from its edge, almost certain to fall into the black hole, feeding its bottomless well. The observations, which will be published in the journal *Nature*, represent the first direct evidence to support the hypothesis that [black holes](#) feed on clouds of [cold gas](#).

The results also suggest that fueling a black hole—a process known as accretion—is a whole lot messier than scientists had once thought.

"The simple model of [black hole accretion](#) consists of a black hole surrounded by a sphere of hot gas, and that gas accretes smoothly onto the black hole, and everything's simple, mathematically," says Michael McDonald, assistant professor of physics in MIT's Kavli Institute for Astrophysics and Space Research. "But this is the most compelling evidence that this process is not smooth, simple, and clean, but actually quite chaotic and clumpy."


Given the new observations, McDonald says black holes probably have two ways of feeding: For most of the time, they may slowly graze on a steady diet of diffuse hot gas. Once in a while, they may quickly gobble up clumps of cold gas as it comes nearby.

"This diffuse, hot gas is available to the black hole at a low level all the time, and you can have a steady trickle of it going in," McDonald says. "Every now and then, you can have a rainstorm with all these droplets of cold gas, and for a short amount of time, the black hole's eating very quickly. So the idea that there are these two dinner modes for black holes is a pretty nice result."

McDonald is a co-author on the paper, which was led by Grant Tremblay, an astronomer at Yale

University.

particular galaxy is among the brightest in the universe, as it is likely producing many new stars.



Deep in the heart of the Abell 2597 Brightest Cluster Galaxy, astronomers see a small cluster of giant gas clouds raining in on the central black hole. They were revealed by the billion light-year-long shadows they cast toward Earth. These ALMA data present the first observational evidence for predicted "chaotic, cold" accretion on a supermassive black hole. Credit: NRAO/AUI/NSF; D. Berry / SkyWorks; ALMA (ESO/NAOJ/NRAO)

Seeing shadows

The researchers made their detection using the Atacama Large Millimeter/submillimeter Array, or ALMA—one of the most powerful telescopes in the world, designed to see the oldest, most distant galaxies in the universe. The team focused ALMA's telescopes 1 billion light years away, on the central galaxy in the Abell 2597 Cluster, a galaxy that is some tens of thousands of [light years](#) across. This


The team originally wanted to get a sense for how many stars this cluster was churning out, so they mapped all the cold gas within the cluster. This cold gas has cooled and condensed out of the diffuse halo of hot gas surrounding a cluster, forming clumps. It is the collapse of cold gas that creates new stars, especially in the cluster's central galaxy.

"In the center of a cluster, there's a single massive galaxy, the big daddy galaxy of the cluster," McDonald says. "It's sitting at the bottom of a gravitational funnel, and all the gas from a thousand galaxies is available to it. These are the galaxies that are the most massive, with the most massive black holes in the universe, and the most potential for star formation."

The researchers used ALMA to map the spectral signatures, or radio emissions, from the galaxy cluster, looking specifically for signatures of carbon monoxide, the presence of which usually indicates very cold gas, of minus 200 degrees Fahrenheit and below. They mapped carbon monoxide across the entire [galaxy cluster](#) and found that as they looked further into the cluster, they encountered progressively cooler gas, from millions of degrees Fahrenheit to subzero temperatures.

At the very center, just at the edge of the cluster's [supermassive black hole](#), the researchers discovered something quite unexpected: the shadows of three very cold, very clumpy gas clouds. The shadows were cast against bright jets of material spewing from the black hole, suggesting that these clouds were very close to being consumed by the black hole.

"We got very lucky," McDonald says. "We could probably look at 100 galaxies like this and not see what we saw just by chance. Seeing three shadows at once is like discovering not just one exoplanet, but three in the first try. Nature was very kind in this case."



"We're only seeing this tiny sliver," McDonald says. "If there are three clouds in just our line of sight, there might be millions of clouds all around. And there's a tremendous amount of energy in just these three [clouds](#). So if we were to look at this thing a million years later, we might see that the black hole is in outburst—much brighter, with more powerful jets, because all this high-energy material is landing on it."

More information: Grant R. Tremblay et al, Cold, clumpy accretion onto an active supermassive black hole, *Nature* (2016). [DOI: 10.1038/nature17969](https://doi.org/10.1038/nature17969)

Provided by Massachusetts Institute of Technology

Composite image of Abell 2597 Brightest Cluster Galaxy. The background image (blue) is from the Hubble Space Telescope. The foreground (red) is ALMA data showing the distribution of carbon monoxide gas in and around the galaxy. The pull-out box is the ALMA data of the "shadow" (black) produced by absorption of the millimeter-wavelength light emitted by electrons whizzing around powerful magnetic fields near the galaxy's supermassive black hole. The shadow indicates that cold clouds of molecular gas are raining in on the black hole. Credit: B. Saxton (NRAO/AUI/NSF); G. Tremblay et al.; NASA/ESA Hubble; ALMA (ESO/NAOJ/NRAO)

A high-energy feast

The team estimated the velocities of the three clouds to be 240, 275, and 355 kilometers per second, with all three headed toward the black hole. McDonald says these three cold gas clouds will likely not stream straight into the black hole but instead be absorbed into its accretion disc—the massive disc of material that will eventually spiral into the black hole.

He adds that while ALMA was only able to see three clouds of cold gas near the black hole, there may be even more in the vicinity, setting the black hole up for quite a feast.

APA citation: Scientists observe supermassive black hole feeding on cold gas (2016, June 8) retrieved 22 April 2021 from <https://phys.org/news/2016-06-scientists-supermassive-black-hole-cold.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.