

# Atrazine alternatives in sweet corn

July 19 2016

---

Atrazine has been very good at killing weeds in corn fields for more than 50 years. But some of the properties that make it a successful herbicide, such as its persistence in the soil and ability to be transported in water, also lead to concerns about potential environmental impact. At both federal and state levels, increasing restrictions on atrazine use has the sweet corn industry wondering about alternatives.

"The list of herbicides available for use on [sweet corn](#) is much more limited than [field corn](#), so historically sweet corn growers and processors have relied heavily on atrazine. They have been in need of alternatives, but the call for those alternatives has gone largely unanswered," says University of Illinois and USDA Agricultural Research Service weed scientist Marty Williams.

Williams and a team of researchers from Oregon and Minnesota polled growers and processors to find out the weed management tactics they would consider using as alternatives.

"At the onset of this project, we engaged the processing sweet corn industry. We knew alternatives to atrazine had to be pragmatic. Are weed biocontrol agents an option? Would a cover crop solve the problem? Is handweeding cost-effective? In this case, no, no, and no. They wanted us to develop systems that integrated mechanical control with existing herbicide technology," Williams recalls.

Most of the alternative herbicides they tried were already registered for use in sweet corn, and came from different herbicide classes than

atrazine. The research team settled on 16 treatments, including three standard atrazine-containing treatments. Treatments were compared in sweet corn fields in Illinois, Minnesota, and Oregon over four years.

"Standard treatments containing atrazine and mesotrione applied postemergence provided the best control, and resulted in crop yields comparable to the weed-free check," Williams states. "In our study, alternative treatments with tembotrione applied postemergence with or without interrow cultivation were similar in weed control and crop yield to atrazine-containing treatments."

Interestingly, interrow cultivation sometimes stimulated weed emergence, thereby reducing effectiveness of the tactic in certain fields. The researchers also found that several other atrazine-free treatments worked well, but only in Oregon.

"Oregon fields were dominated by small-seeded weed species that were controlled by more products than the large-seeded weeds more common in the Midwest, such as velvetleaf and giant ragweed," Williams explains.

Tembotriione was used to represent the HPPD inhibiting class of herbicides, which includes other efficacious herbicides such as topramezone, mesotrione, and bicyclopyrone. This mode of action was developed most recently, but some weeds are already becoming resistant to HPPD inhibitors.

"As with most herbicides, misuse of the alternatives to atrazine could select for herbicide resistant weed populations. Nonetheless, it is possible to grow sweet corn without atrazine using an approach that sweet corn growers and processors consider adoptable," Williams notes.

**More information:** The article, "Alternatives to atrazine for weed

management in processing sweet corn," is published in *Weed Science*.

Provided by University of Illinois at Urbana-Champaign

Citation: Atrazine alternatives in sweet corn (2016, July 19) retrieved 20 September 2024 from <https://phys.org/news/2016-07-atrazine-alternatives-sweet-corn.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.