

Towards a better screen: New molecules promise cheaper, more efficient OLED displays

8 August 2016



Alán Aspuru-Guzik, Professor of Chemistry and Chemical Biology at Harvard University. Credit: Stephanie Mitchell

Harvard University researchers have designed more than 1,000 new blue-light emitting molecules for organic light-emitting diodes (OLEDs) that could dramatically improve displays for televisions, phones, tablets and more.

OLED screens use [organic molecules](#) that emit light when an electric current is applied. Unlike ubiquitous liquid crystal displays (LCDs), OLED screens don't require a backlight, meaning the display can be as thin and flexible as a sheet of plastic. Individual pixels can be switched on or entirely off, dramatically improving the screen's color contrast and energy consumption. OLEDs are already replacing LCDs in high-end consumer devices but a lack of stable and efficient blue materials has made them less competitive in large displays such as televisions.

The interdisciplinary team of Harvard researchers, in collaboration with MIT and Samsung, developed

a large-scale, computer-driven screening process, called the Molecular Space Shuttle, that incorporates theoretical and experimental chemistry, [machine learning](#) and cheminformatics to quickly identify new OLED molecules that perform as well as, or better than, industry standards.

"People once believed that this family of organic light-emitting molecules was restricted to a small region of molecular space," said Alán Aspuru-Guzik, Professor of Chemistry and Chemical Biology, who led the research. "But by developing a sophisticated molecular builder, using state-of-the-art machine learning, and drawing on the expertise of experimentalists, we discovered a large set of high-performing blue OLED materials."

The research is described in the current issue of *Nature Materials*.

The biggest challenge in manufacturing affordable OLEDs is emission of the color blue.

Like LCDs, OLEDs rely on green, red and blue subpixels to produce every color on screen. But it has been difficult to find organic molecules that efficiently emit blue light. To improve efficiency, OLED producers have created organometallic molecules with expensive transition metals like iridium to enhance the molecule through phosphorescence. This solution is expensive and it has yet to achieve a stable blue color.

Aspuru-Guzik and his team sought to replace these organometallic systems with entirely organic molecules.

The team began by building libraries of more than 1.6 million candidate molecules. Then, to narrow the field, a team of researchers from the Harvard

John A. Paulson School of Engineering and Applied Sciences (SEAS), led by Ryan Adams, Assistant Professor of Computer Science, developed new machine learning algorithms to predict which molecules were likely to have good outcomes, and prioritize those to be virtually tested. This effectively reduced the computational cost of the search by at least a factor of ten.

To help bridge the gap between theoretical modeling and experimental practice, Hirzel and the team built a web application for collaborators to explore the results of more than half a million quantum chemistry simulations.

Every month, Gómez-Bombarelli and coauthor Jorge Aguilera-Iparraguirre, also a [postdoctoral fellow](#) in the Aspuru-Guzik lab, selected the most promising molecules and used their software to create "baseball cards," profiles containing important information about each molecule.

This process identified 2500 molecules worth a closer look. The team's experimental collaborators at Samsung and MIT then voted on which molecules were most promising for application. The team nicknamed the voting tool "molecular Tinder" after the popular online dating app.

"We facilitated the social aspect of the science in a very deliberate way," said Hirzel.

"The computer models do a lot but the spark of genius is still coming from people," said Gómez-Bombarelli.

"The success of this effort stems from its multidisciplinary nature," said Aspuru-Guzik. "Our collaborators at MIT and Samsung provided critical feedback regarding the requirements for the molecular structures."

"The high throughput screening technique pioneered by the Harvard team significantly reduced the need for synthesis, experimental characterization, and optimization," said Marc Baldo, Professor of Electrical Engineering and Computer Science at MIT and coauthor of the paper. "It shows the industry how to advance OLED technology faster and more efficiently."

After this accelerated design cycle, the team was left with hundreds of molecules that perform as well as, if not better than, state-of-the-art metal-free OLEDs.

Applications of this type of molecular screening

also extend far beyond OLEDs.

"This research is an intermediate stop in a trajectory towards more and more advanced organic [molecules](#) that could be used in flow batteries, solar cells, organic lasers, and more," said Aspuru-Guzik. "The future of accelerated molecular design is really, really exciting."

More information: Design of efficient molecular organic light-emitting diodes by a high-throughput virtual screening and experimental approach, *Nature Materials*, [DOI: 10.1038/nmat4717](https://doi.org/10.1038/nmat4717)

Provided by Harvard John A. Paulson School of Engineering and Applied Sciences

APA citation: Towards a better screen: New molecules promise cheaper, more efficient OLED displays (2016, August 8) retrieved 6 May 2021 from <https://phys.org/news/2016-08-screen-molecules-cheaper-efficient-oled.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.