

Logging can decrease water infiltration into forest soils, study finds

August 17 2016



Langston Simmons, former MU graduate student and lead author on the study, takes soil samples from a logging forest. Credit: MU News Bureau

Soil water infiltration, or the ability of soil to absorb water and allow it to move through different soil layers, is an important environmental factor in forests, especially forests undergoing logging operations. This property can affect how quickly those forests can regenerate after being logged. Now, University of Missouri researchers have found that logging operations can negatively affect soil density and water infiltration within forests, particularly along makeshift logging roads and landing areas where logs are stored before being trucked to sawmills. Stephen Anderson, the William A. Albrecht Distinguished Professor of Soil Science at Mizzou, says changing the soil density and water infiltration within forests can cause many different problems.

"We found that along these logging roads and landing areas, the soil was more dense and compact with slower water infiltration than in the surrounding, untouched areas of the [forest](#)," Anderson said. "This can cause many environmental challenges in forests because dense soil prevents rainwater from soaking in; rather, this water will run off and cause erosion. This erosion can carry fertile topsoil away from forests, which enters streams and makes it difficult for those forests being logged to regenerate with new growth as well as polluting surface water resources."

For their study, Anderson and his former graduate student, Langston Simmons, took soil core samples up to 40 centimeters deep from logging roads, log landing areas and logged areas in portions of the Mark Twain National Forest in Callaway County, Missouri. They found that the soil from logging roads and landing areas was more dense, had much slower water infiltration and lower water retention capacity than the areas of forest that had been logged. Anderson says this study shows the need for treatment of these impacted areas within logged forests.



Soil samples being analyzed in an MU laboratory. Credit: MU News Bureau

"It is clear that even though logging companies can take precautions to prevent many types of negative environmental impacts from their operations, soil density and [water infiltration](#) are being negatively affected," Anderson said. "It is important these areas of compacted soil be identified and treated to reduce soil compaction and prevent long-term effects on forest regeneration and production. It is in the land managers' best interests to ensure that forest soils remain a healthy density because dense soil can lead to reduced tree production and poor wood quality for future logging operations."

This study, "Effects of logging activities on selected [soil](#) physical and

hydraulic properties for a claypan landscape," was published in *Geoderma*.

Provided by University of Missouri-Columbia

Citation: Logging can decrease water infiltration into forest soils, study finds (2016, August 17) retrieved 19 September 2024 from

<https://phys.org/news/2016-08-decrease-infiltration-forest-soils.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.