


# Scientists present the most accurate time measurements of quantum jumps to date

7 November 2016



A laser pulse hits a helium atom. One of the electron is ripped out of the atom, the other electron may change its quantum state. Credit: TU Wien

When a quantum system changes its state, this is called a quantum jump. Usually, these quantum jumps are considered to be instantaneous. Now, new methods for high-precision measurements allow us to study the time evolution of these quantum jumps. On a time scale of attoseconds, their time structure becomes visible. It is the most accurate time measurement of quantum jumps to date.

Quantum particles can change their state very quickly—this is called a "quantum jump". An atom, for example, can absorb a photon, thereby changing into a state of higher energy. Usually, these processes are thought to happen instantaneously, from one moment to the next. However, with new methods, developed at TU Wien (Vienna), it is now possible to study the time structure of such extremely fast state changes. Much like an electron microscope allows us to take a look at tiny structures which are too small to be


seen with the naked eye, [ultrashort laser pulses](#) allow us to analyse temporal structures which used to be inaccessible.

The theoretical part of the project was done by Prof. Joachim Burgdörfer's team at TU Wien (Vienna), which also developed the initial idea for the experiment. The experiment was performed at the Max-Planck-Institute for Quantum Optics in Garching (Germany). The results have now been published in the journal *Nature Physics*.

## The Most Accurate Time Measurement of Quantum Jumps

A neutral helium atom has two electrons. When it is hit by a high energy laser pulse, it can be ionized: one of the electrons is ripped out of the atom and departs from it. This process occurs on a time scale of attoseconds - one attosecond is a billionth of a billionth of a second.

"One could imagine that the other electron, which stays in the atom, does not really play an important part in this process—but that's not true", says Renate Pazourek (TU Wien). The two electrons are correlated, they are closely connected by the laws of quantum physics, they cannot be seen as independent particles. "When one electron is removed from the atom, some of the laser energy can be transferred to the second electron. It remains in the atom, but it is lifted up to a state of higher energy", says Stefan Nagele (TU Wien).



"instantaneous" can now be seen as temporal developments which can be calculated, measured and even controlled. This does not only help to understand the basic laws of nature, it also brings new possibilities of manipulating matter on a quantum scale.

**More information:** Attosecond correlation dynamics, *Nature Physics*, [nature.com/articles/doi:10.1038/nphys3941](http://nature.com/articles/doi:10.1038/nphys3941)

Provided by Vienna University of Technology

Once a photon has removed an electron from a helium atom, it is possible to calculate the probable position of the remaining electron. The likeliest position of the electron is shown in the image as the brightest area around the atomic nucleus (which itself is not visible in the image) Credit: M. Ossiander (TUM) / M. Schultz (MPQ)

Therefore, it is possible to distinguish between two different ionization processes: one, in which the remaining electron gains additional energy and one, in which it remains in a state of minimal energy. Using a sophisticated experimental setup, it was possible to show that the duration of these two processes is not exactly the same.

"When the remaining electron jumps to an excited state, the photo ionization process is slightly faster - by about five attoseconds", says Stefan Nagele. It is remarkable how well the experimental results agree with theoretical calculations and large-scale computer simulations carried out at the Vienna Scientific Cluster, Austria's largest supercomputer: "The precision of the experiment is better than one attosecond. This is the most accurate time measurement of a [quantum jump](#) to date", says Renate Pazourek.

### Controlling Attoseconds

The experiment provides new insights into the physics of ultrashort time scales. Effects, which a few decades ago were still considered

APA citation: Scientists present the most accurate time measurements of quantum jumps to date (2016, November 7) retrieved 19 June 2021 from <https://phys.org/news/2016-11-scientists-accurate-quantum-date.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*