


# Second-generation stars identified, giving clues about their predecessors

6 December 2016, by Brian Wallheimer



The figure shows a sub-population of ancient stars, called Carbon-Enhanced Metal-Poor (CEMP) stars. These stars contain 100 to 1,000,000 times LESS iron (and other heavy elements) than the Sun, but 10 to 10,000 times MORE carbon, relative to iron. The unusual chemical compositions of these stars provides clues to their birth environments, and the nature of the stars in which the carbon formed. In the figure, A(C) is the absolute amount of carbon, while the horizontal axis represents the ratio of iron, relative to hydrogen, compared with the same ratio in the Sun. Credit: University of Notre Dame

University of Notre Dame astronomers have identified what they believe to be the second generation of stars, shedding light on the nature of the universe's first stars.

A subclass of carbon-enhanced metal-poor (CEMP) [stars](#), the so-called CEMP-no stars, are ancient stars that have large amounts of carbon but little of the [heavy metals](#) (such as iron) common to later-generation stars. Massive first-generation stars made up of pure hydrogen and helium produced and ejected [heavier elements](#) by

stellar winds during their lifetimes or when they exploded as supernovae. Those metals—anything heavier than helium, in astronomical parlance—polluted the nearby [gas clouds](#) from which new stars formed.

Jinmi Yoon, a postdoctoral research associate in the Department of Physics; Timothy Beers, the Notre Dame Chair in Astrophysics; and Vinicius Placco, a research professor at Notre Dame, along with their collaborators, show in findings published in the *Astrophysics Journal* this week that the lowest metallicity stars, the most chemically primitive, include large fractions of CEMP stars. The CEMP-no stars, which are also rich in nitrogen and oxygen, are likely the stars born out of hydrogen and helium gas clouds that were polluted by the elements produced by the universe's first stars.

"The CEMP-no stars we see today, at least many of them, were born shortly after the Big Bang, 13.5 billion years ago, out of almost completely unpolluted material," Yoon says. "These stars, located in the halo system of our galaxy, are true second-generation stars—born out of the nucleosynthesis products of the very first stars."

Beers says it's unlikely that any of the universe's first stars still exist, but much can be learned about them from detailed studies of the next generation of stars.

"We're analyzing the chemical products of the very first stars by looking at what was locked up by the second-generation stars," Beers says. "We can use this information to tell the story of how the first elements were formed, and determine the distribution of the masses of those first stars. If we know how their masses were distributed, we can model the process of how the first stars formed and evolved from the very beginning."

The authors used high-resolution spectroscopic

data gathered by many astronomers to measure the chemical compositions of about 300 stars in the halo of the Milky Way. More and heavier elements form as later generations of stars continue to contribute additional metals, they say. As new generations of stars are born, they incorporate the metals produced by prior generations. Hence, the more heavy metals a star contains, the more recently it was born. Our sun, for example, is relatively young, with an age of only 4.5 billion years.

A companion paper, titled "Observational constraints on first-star nucleosynthesis. II. Spectroscopy of an ultra metal-poor CEMP-no star," of which Placco was the lead author, was also published in the same issue of the journal this week. The paper compares theoretical predictions for the chemical composition of zero-metallicity supernova models with a newly discovered CEMP-no star in the Milky Way galaxy.

**More information:** Jinmi Yoon et al.

OBSERVATIONAL CONSTRAINTS ON FIRST-STAR NUCLEOSYNTHESIS. I. EVIDENCE FOR MULTIPLE PROGENITORS OF CEMP-NO STARS, *The Astrophysical Journal* (2016). [DOI: 10.3847/0004-637X/833/1/20](https://doi.org/10.3847/0004-637X/833/1/20)

Vinicius M. Placco et al. OBSERVATIONAL CONSTRAINTS ON FIRST-STAR NUCLEOSYNTHESIS. II. SPECTROSCOPY OF AN ULTRA METAL-POOR CEMP-no STAR, *The Astrophysical Journal* (2016). [DOI: 10.3847/0004-637X/833/1/21](https://doi.org/10.3847/0004-637X/833/1/21)

Provided by University of Notre Dame

APA citation: Second-generation stars identified, giving clues about their predecessors (2016, December 6) retrieved 24 February 2021 from <https://phys.org/news/2016-12-second-generation-stars-clues-predecessors.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*