


# Can the donut-shaped magnet 'CAPPuccino submarine' hunt for dark matter?

January 23 2017

---



Scientists at IBS CAPP are prototyping haloscopes - machines that hunt for dark matter. Haloscope have very strong magnets. Helix-shaped magnets (solenoid magnets, on the left) are commonly used in dark matter experiments. CAPP scientists are also investigating the possibility of using donut-shaped magnets, technically known as toroidal magnets, and nicknamed this device the CAPPuccino submarine. Credit: Institute for Basic Science

Although it sounds hard to believe, everything we see with the naked eye or through microscopes and telescopes accounts for just 4 percent of the known universe. The rest comprises dark energy (69 percent) and dark matter (27 percent). Although there seems to be more dark matter than visible matter in the universe, we still have not been able to directly detect it. The reason is that dark matter does not emit light or absorb electromagnetic waves, so it is really hard to observe. Interestingly, dark matter is needed to explain the motions of galaxies and some of the current theories of galaxy formation and evolution. For example, the galaxy that contains our solar system, the Milky Way, seems to be enveloped by a much larger halo of dark matter; though invisible, its existence is inferred through its effects on the motions of stars and gases.

Although [dark matter](#) particles have not been detected so far, scientists know that these particles have a very small mass and are distributed throughout the universe. One [dark matter particle](#) candidate is the axion. Axions have extremely weak interactions with matter and so scientists need special equipment to catch their presence. Specifically, scientists use the so-called axion to two-photons coupling technique, which takes advantage of the fact that an axion passing through a strong magnetic field can interact with a photon and convert into another photon. To record this interaction, IBS scientists are in the process of building haloscopes in Daejeon in South Korea.

Haloscopes contain resonant cavities immersed in an extra-strong magnetic field. "In simple terms, you can image the resonant cavity as a cylinder, like a soft drink can, where the energy of the photons generated from the axions-photons interaction is amplified," explains KO Byeong Rok, first author of this study.



Artistic impression of the Milky Way galaxy with the mysterious dark matter halo shown in blue, but actually invisible. Credit: ESO/L. Calçada, Wikipedia

The magnets used for these types of experiments have the shape of a coil wound into a helix, technically known as a solenoid. However, depending on the height of the magnet, there is the risk of losing the signal coming from the axion-photon interaction. For this reason, IBS scientists decided to look deeper into another type of magnets shaped like donuts, called toroidal magnets.

"Magnets are the most important feature of the haloscope, and also the most expensive. While other experiments seeking to detect dark matter around the world use solenoid magnets, we are the first to try to use toroidal magnets. Since they have never been used before, you cannot easily buy the equipment, so we develop it ourselves," explains Professor Ko.

In order to hunt the axion, scientists need to get out in front of it, and predict the magnitude of the [electromagnetic energy](#) expected from the axion-to-photon conversion. Electromagnetic energy is the sum of electric and magnetic energies. Both can be easily calculated for a solenoid magnet, but if the magnet is toroidal, it is practically impossible to calculate the [magnetic energy](#) analytically. Because of this, it was believed that toroidal magnets could not be used for the haloscope.

This paper from IBS shows the opposite. Starting from an adjusted version of the Maxwell equation, which defines how charged particles give rise to electric and magnetic forces, scientists found that electric energy and magnetic energy from the axion-photon interaction are equal in both types of magnets. Therefore, even though the magnetic energy of a toroidal magnet is unknown, in order to obtain the electromagnetic energy which is the sum of the two, it is possible to double up the electric energy and obtain the magnetic energy.

Another finding is that the energy emitted from the interaction and conversion of the axion to photon is independent from the position of the cavity inside a solenoid magnet. However, this is not the case for toroid magnets.

IBS CAPP scientists have nicknamed the toroidal cavity "CAPPuccino submarine" because its color resembles the beverage, and its particular shape. All the theoretical findings published in this paper are going to form a solid background for the development and prototyping of new machines for the search of dark matter.

**More information:** B. R. Ko et al. Electric and magnetic energy at axion haloscopes, *Physical Review D* (2016). [DOI: 10.1103/PhysRevD.94.111702](https://doi.org/10.1103/PhysRevD.94.111702)

Provided by Institute for Basic Science

Citation: Can the donut-shaped magnet 'CAPPuccino submarine' hunt for dark matter? (2017, January 23) retrieved 19 September 2024 from <https://phys.org/news/2017-01-donut-shaped-magnet-cappuccino-submarine-dark.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.