


Astronomers uncover properties of a molecular outflow feature in a nearby starburst galaxy

24 January 2017, by Tomasz Nowakowski



ALMA CO(1–0) observations of NGC 253. Credit: Walter et al., 2017.

(Phys.org)—An international team of astronomers led by Fabian Walter of the Max Planck Institute for Astronomy in Germany has revealed essential properties of a molecular outflow feature in the nearby starburst galaxy designated NGC253. The findings were presented in a paper published Jan. 18 on the arXiv pre-print repository.

NGC 253, also known as the Sculptor Galaxy, is an intermediate spiral starburst galaxy in the constellation Sculptor some 11 million [light years](#) away. The galaxy is one of the brightest spiral [galaxies](#) visible and also one of the dustiest. Due to its proximity and the fact that it is currently undergoing a period of intense star formation, it serves as one of the best laboratories to study starburst–driven galactic–scale winds in detail.

NGC 253 is known for the galactic wind emerging from its central area, which carries significant amounts of [molecular gas](#). In order to better characterize this emission, Walter and his team carried out a series of observations between November 2013 and August 2014 utilizing the Atacama Large Millimeter/submillimeter Array (ALMA) in Chile. This observational campaign

allowed the researchers to uncover crucial properties of the studied molecular outflow.

"We present a detailed study of a molecular outflow feature in the nearby starburst galaxy NGC 253 using ALMA," the paper reads.

According to the study, NGC 253's most prominent outflow feature towards the south has a projected length of about 978 light years, a width of approximately 163 light years, and a velocity dispersion of 40 km s^{-1} . The scientists note that its extent and dispersion are consistent with an ejection from the galaxy's disk starting about 1 million years ago.

The kinematics of the molecular gas in this feature show that the material is approaching us and is also accelerating. However, the astronomers were unable to determine whether or not the molecular mass entrained in the outflow will escape the galaxy, or be recycled to fuel later episodes of [star formation](#). More observations are required to draw final conclusions.

"The kinematics of the molecular gas are consistent with accelerating with a velocity gradient of $1 \text{ km s}^{-1} \text{ pc}^{-1}$, and at its last measurable point it approaches the escape velocity. (...) More sensitive ALMA observations will trace the outflow even further out, and will shed light on whether or not part of the ejected molecular material will escape the galaxy," the researchers wrote in the paper.

The team also discovered bright tracers of dense molecular gas like HCN, CN, HCO+, CS in the molecular outflow. Moreover, they found that line ratios HCN/CO measured in the outflow are high and consistent with ratios observed in the central starburst region of NGC253 and in other starbursts, while the HCN/CO line ratio in the galaxy's disk is

significantly lower, which is typical for gas in the disks of nearby galaxies. The researchers concluded that this indicates that the dense molecular gas is retaining its properties in the process of its ejection from the central regions into the outflow.

More information: Dense Molecular Gas Tracers in the Outflow of the Starburst Galaxy NGC 253, arXiv:1701.05040 [astro-ph.GA]
arxiv.org/abs/1701.05040

Abstract

We present a detailed study of a molecular outflow feature in the nearby starburst galaxy NGC 253 using ALMA. We find that this feature is clearly associated with the edge of NGC 253's prominent ionized outflow, has a projected length of ~ 300 pc, with a width of ~ 50 pc and a velocity dispersion of ~ 40 km s⁻¹, consistent with an ejection from the disk about 1 Myr ago. The kinematics of the molecular gas in this feature can be interpreted (albeit not uniquely) as accelerating at a rate of 1 km s⁻¹ pc⁻¹. In this scenario, the gas is approaching escape velocity at the last measured point. Strikingly, bright tracers of dense molecular gas (HCN, CN, HCO⁺, CS) are also detected in the molecular outflow: We measure an HCN(1-0)/CO(1-0) line ratio of $\sim 1/10$ in the outflow, similar to that in the central starburst region of NGC 253 and other starburst galaxies. By contrast, the HCN/CO line ratio in the NGC 253 disk is significantly lower ($\sim 1/30$), similar to other nearby galaxy disks. This strongly suggests that the streamer gas originates from the starburst, and that its physical state does not change significantly over timescales of ~ 1 Myr during its entrainment in the outflow. Simple calculations indicate that radiation pressure is not the main mechanism for driving the outflow. The presence of such dense material in molecular outflows needs to be accounted for in simulations of galactic outflows.

© 2017 Phys.org

APA citation: Astronomers uncover properties of a molecular outflow feature in a nearby starburst galaxy (2017, January 24) retrieved 20 January 2021 from <https://phys.org/news/2017-01-astronomers-uncover-properties-molecular-outflow.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.