

Coal mine dust hastens Arctic snow melt

February 1 2017



Collecting spectral reflectance measurements of surface snow with corresponding snow samples in Northern Svalbard. Credit: Alia Khan / University of Colorado Boulder

Dust released by an active coal mine in Svalbard, Norway, reduced the spectral reflectance of nearby snow and ice by up to 84 percent, according to new University of Colorado Boulder-led research.

The study illustrates the significant, localized role that dark-colored particulates—which absorb more solar radiation than light-colored snow and keep more heat closer to the Earth's surface—can play in hastening Arctic ice melt.

The study was published today in the *Journal of Geophysical Research: Atmospheres*, a publication of the American Geophysical Union.

Unblemished snow and ice have a very high spectral albedo, a measurement used to indicate how effectively a given surface reflects solar energy. Over time, airborne black carbon particles (from soot or automobile emissions, for example) or other mineral dust can travel long distances in the atmosphere and settle on snow and glaciers, lowering the overall albedo.

To study the localized effects of coal dust on an area with high spectral reflectance, CU Boulder researchers focused on an active coal mine in Svalbard, Norway, located on a sparsely populated island north of the Arctic Circle. The researchers collected snow and ice samples from four sites at varying distances from the mine, with some samples being visibly dirtier than others.



Collecting spectral reflectance measurements of surface snow with corresponding snow samples next to Mine 7 near Longyearbyen, Svalbard. Credit: Alia Khan / University of Colorado Boulder

The researchers then measured the light absorption capacity of each sample, adjusting for environmental factors such as snow grain size and location relative to the mine. Overall, the study concluded that coal dust had a strong but localized effect, reducing the albedo in the immediate area by up to 84 percent.

The findings may provide a foundation for similar research using satellites and remote sensing techniques in far-flung areas.

"The extreme contrast between [snow](#) and dust at this particular site gave

us a baseline to develop algorithms that we can now use to take future measurements in areas that aren't easily accessible," said lead study author Alia Khan, a post-doctoral researcher in CU Boulder's National Snow and Ice Data Center and former graduate student at the Institute of Arctic and Alpine Research (INSTAAR).

The findings may also provide context for future policymaking decisions regarding the potential expansion of mining the coal-rich Arctic region, especially in light of ongoing permafrost thaw that may allow more land-based drilling operations.

"We hope these ground-based spectral measurements could be used in the management of future energy development in the Arctic, especially for mines that may be unavailable for ground-based observations, but may be large enough to be visible by satellite," said Khan.

More information: Impacts of coal dust from an active mine on the spectral reflectance of Arctic surface snow in Svalbard, Norway, *Journal of Geophysical Research: Atmospheres*, [DOI: 10.1002/2016JD025757](https://doi.org/10.1002/2016JD025757) , [onlinelibrary.wiley.com/doi/10 ... 02/2016JD025757/full](https://onlinelibrary.wiley.com/doi/10.1002/2016JD025757/full)

Provided by University of Colorado at Boulder

Citation: Coal mine dust hastens Arctic snow melt (2017, February 1) retrieved 19 April 2024 from <https://phys.org/news/2017-02-coal-hastens-arctic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.