


# New algorithm identifies gene transfers between different bacterial species

February 24 2017



Mosaic pneumococcal population structure caused by horizontal gene transfer is shown on the left for a subset of genes. Matrix on the right shows a genome-wide summary of the relationships between the bacteria, ranging from blue (distant) to yellow (closely related). Credit: Pekka Marttinen

Gene transfers are particularly common in the antibiotic-resistance genes of *Streptococcus pneumoniae* bacteria.

When mammals breed, the genome of the offspring is a combination of the parents' genomes. Bacteria, by contrast, reproduce through cell division. In theory, this means that the genomes of the offspring are copies of the parent genome. However, the process is not quite as straightforward as this due to horizontal [gene transfer](#) through which bacteria can transfer fragments of their genome to each other. As a result of this phenomenon, the genome of an individual bacterium can be a combination of genes from several different donors. Some of the genome fragments may even originate from completely different species.

In a recent study combining machine learning and bioinformatics, a new computational method was developed for modelling gene transfers between different lineages of a bacterial population or even between entirely different bacterial species. The method was used to analyse a collection of 616 whole-genomes of a recombinogenic pathogen *Streptococcus pneumoniae*.

## **The usefulness of a gene affects its transfer rate**

In the study, several individual genes in which gene transfers were considered particularly common were identified. These genes also included genes causing resistance to antibiotics.

'In the case of antibiotic-resistance genes, the number of gene transfers may be related to how useful these genes are to bacteria and to the resulting selection pressure', says Academy Research Fellow Pekka Marttinen from the Aalto University Department of Computer Science.

'The study will not provide a direct solution to antibiotic resistance because this would require a profound understanding of how the resistance occurs and spreads. Nevertheless, knowing the extent to which gene transfer occurs between different species and lineages can help in

improving this understanding', he explains.

The study was able to show that gene transfer occurs both within species and between several different species. The large number of transfers identified during the study was a surprise to the researchers.

'Previous studies have shown that gene transfers are common in individual [genes](#), but our team was the first to use a computational method to show the extent of gene transfer across the entire [genome](#)', Marttinen says.

'The method also makes it possible to effectively examine ancestral gene transfers for the first time, which is important in examining transfers between different [species](#).'

*Molecular Biology and Evolution* published the results in February.

**More information:** Rafal Mostowy et al, Efficient inference of recent and ancestral recombination within bacterial populations, *Molecular Biology and Evolution* (2017). [DOI: 10.1093/molbev/msx066](https://doi.org/10.1093/molbev/msx066)

Provided by Aalto University

Citation: New algorithm identifies gene transfers between different bacterial species (2017, February 24) retrieved 24 April 2024 from <https://phys.org/news/2017-02-algorithm-gene-bacterial-species.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.