

Reading between the lines of highly turbulent plasmas

29 March 2017

Plasma, the ionised state of matter found in stars, is still not fully understood, largely due to its instability. Astrophysicists have long sought to develop models that can account for the turbulent motions inside plasma, based on observing line shapes emitted by atoms and ions in the plasma. Turbulences are typically detected through the observation of broadened lines due to the Doppler effect, similar to the principle behind radar.

In a new study published in *EPJ D*, Roland Stamm from the CNRS and Aix-Marseille University, France, and colleagues develop an iterative simulation model that accurately predicts, for the first time, the changes to the line [shape](#) in the presence of strong plasma [turbulence](#).

Ultimately, the authors aim to provide a system for assessing plasma turbulence that is valid for both a stellar atmosphere and the ITER tokamak designed to generate fusion energy. Line shapes are extensively employed as a powerful diagnostic tool for detecting turbulences in stable gases and plasmas. For many years now, astrophysicists have developed and employed models that gauge the [effect](#) of turbulent motions in the broadening of line shapes due to the Doppler effect. Such models are now also being employed to understand the role of turbulences in plasmas created to harvest energy from fusion.

In this study, the authors review the effects of strong turbulence on the line shapes when the plasma is subjected to an external energy source, such as a beam of charged particles. Their model accounts for the effect of an electric field on a hydrogen atom subjected to strong turbulence within a plasma. They subsequently perform numerical simulations for various low-density plasmas, ultimately determining that the width of the hydrogen line increases in the presence of strong turbulence connected to the external energy source, shaped as a sequence of solitons. Under such conditions, the line shapes show the

presence of waves oscillating at the [plasma](#) frequency. Electrostatic waves experience a cycle during which they rise to very high intensities before dissipating and reforming, drawing [energy](#) from the driver beam.

More information: Roland Stamm et al, Line shapes in turbulent plasmas, *The European Physical Journal D* (2017). [DOI: 10.1140/epjd/e2017-70737-2](#)

Provided by Springer

APA citation: Reading between the lines of highly turbulent plasmas (2017, March 29) retrieved 15 January 2021 from <https://phys.org/news/2017-03-lines-highly-turbulent-plasmas.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.