

Solving the mystery of the Arctic's green ice

March 29 2017



Melt ponds on the surface of Arctic ice. Credit: NASA

In 2011, researchers observed something that should be impossible—a massive bloom of phytoplankton growing under Arctic sea ice in conditions that should have been far too dark for anything requiring photosynthesis to survive. So, how was this bloom possible?


Using mathematical modeling, researchers from the Harvard John A. Paulson School of Engineering and Applied Sciences (SEAS) found that thinning Arctic sea ice may be responsible for these blooms and more blooms in the future, potentially causing significant disruption in the Arctic food chain.

The research is described in *Science Advances* and is a collaboration between researchers from SEAS, University of Oxford and University of Reading.

Phytoplankton underpins the entire Arctic food web. Every summer, when the sea ice retreats, sunlight hitting the open water triggers a massive bloom of plankton. These plumes attract fish, which attract larger predators and provides food for indigenous communities living in the Arctic.

Phytoplankton shouldn't be able to grow under the ice because ice reflects most sunlight light back into space, blocking it from reaching the water below.

But over the past decades, Arctic ice has gotten darker and thinner due to warming temperatures, allowing more and more sunlight to penetrate to the water beneath. Large, dark pools of water on the surface of the ice, known as [melt ponds](#), have increased, lowering the reflectivity of the ice. The ice that remains frozen is thin and getting thinner.



Areas of the Arctic Ocean capable of permitting blooms of phytoplankton under sea ice in July. Credit: Christopher Horvat

"Our big question was, how much sunlight gets transmitted through the sea ice, both as a function of thickness, which has been decreasing, and the melt pond percentage, which has been increasing," said Chris Horvat, first author of the paper and graduate student in applied mathematics at SEAS. "What we found was that we went from a state where there wasn't any potential for plankton blooms to massive regions of the Arctic being susceptible to these types of growth."

The team's mathematical modeling found that while the melt ponds contribute to conditions friendly to blooms, the biggest culprit is ice thickness.

Twenty years ago, only about 3 to 4 percent of Arctic sea ice was thin enough to allow large colonies of plankton to bloom underneath. Today, the researchers found that nearly 30 percent of the ice-covered Arctic Ocean permits sub-ice blooms in summer months.

"The meter decline in [sea ice thickness](#) in the Arctic in the past 30 years has dramatically changed the ecology in that area," said Horvat. "All of a

sudden, our entire idea about how this ecosystem works is different. The foundation of the Arctic food web is now growing at a different time and in places that are less accessible to animals that need oxygen."

The researchers hope their model will be helpful for planning future expeditions to observe these blooms and measuring the impact this shift will have on ecosystems.

More information: "The frequency and extent of sub-ice phytoplankton blooms in the Arctic Ocean," *Science Advances*, advances.sciencemag.org/content/3/3/e1601191

Provided by Harvard John A. Paulson School of Engineering and Applied Sciences

Citation: Solving the mystery of the Arctic's green ice (2017, March 29) retrieved 20 September 2024 from <https://phys.org/news/2017-03-mystery-arctic-green-ice.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.