

More large, high-intensity forest fires likely in coming years

April 6 2017, by Christie Delfanian



This prescribed burn in the Kootenai National Forest in northwestern Montana, is designed to reduce the chances of a large wildfire erupting. However, wildfire experts predict that the number of large, high-intensity fire events will increase, in part due to a 20 to 50 percent increase in the number of days when conditions are conducive to fires. Credit: South Dakota State University

When it comes to large, high-intensity forest fires, we can expect to see a lot more in the coming years, according to South Dakota State University professor Mark Cochrane, a senior scientist at the Geospatial Sciences Center of Excellence.

Using satellite data from 2002 to 2013, Cochrane and researchers from the University of Tasmania and the University of Idaho examined nearly 23,000 fires worldwide, identifying 478 large, high-intensity fires which they defined as extreme wildfire events. Their work is described in the Feb. 2017 issue of *Nature Ecology and Evolution*.

"Almost all happened under bad conditions—high temperatures, dry conditions and strong winds, which tell us that weather and climate are very important," Cochrane said. Using monthly world weather data from 2000 to 2014, the researchers modeled the likely changes in fire behavior from 2041 to 2070, predicting a 20 to 50 percent increase in the number of days when conditions are conducive to fires.

"Those conditions are based on business-as-usual [carbon emissions](#)," Cochrane continued. "This will continue to worsen after 2070 unless we get very serious about cutting global carbon emissions."

By 2041, there will likely be 35 percent more of these large, [catastrophic fires](#) per decade, according to Cochrane. "That translates to four extreme fire events for every three that occur now."

However, that risk is not spread evenly, Cochrane explained. Forests in the western United States, southeastern Australia, Europe and the eastern Mediterranean region that extends from Greece to Lebanon and Syria are among those areas at highest risk.

Defining extreme fire events

Though the concept of huge, devastating wildfires, sometimes called megafires, has been tossed around, Cochrane said, "There is no operational definition." Therefore, the research team, led by University of Tasmania professor David Bowman, examined fire intensity and area.

First, the researchers identified hotspots using moderate resolution imaging spectroradiometer, or MODIS, from two earth-imaging satellites to measure the amount of heat energy released, known as fire radiative power. To do this, they looked at the total energy being released in each 25,000-acre block across the planet, Cochrane explained. "It's a combination of the area that is burning and the intensity at which it is burning."

Through that analysis, the researchers identified 478 extreme fire events. "We limited ourselves to the top .003 percent," he said. "Anyone would agree that these are pretty intense, large events."

Determining impact on humans

They further narrowed the extreme fire events to ones that had the greatest impact on humans. "Fires in the boreal forest might be very large, but they do not impact many people," he pointed out. "We looked for those in which people had to clear out of the way."

The researchers identified 144 fires that were catastrophic, meaning

people died and homes were destroyed. "Most of these fires were in the western United States and southeastern Australia, which have fairly high population densities," Cochrane said.

Wind-driven fires accounted for nearly 35 percent of these catastrophic events, while severe drought was a factor in nearly 22 percent. Other extreme [fire](#) weather conditions, largely due to high temperatures and low humidity, accounted for slightly more than 20 percent of these costly fires.

"Not only is climate making things worse, but people are building homes in these flammable landscapes," he said.

More information: David M. J. S. Bowman et al. Human exposure and sensitivity to globally extreme wildfire events, *Nature Ecology & Evolution* (2017). [DOI: 10.1038/s41559-016-0058](https://doi.org/10.1038/s41559-016-0058)

Provided by South Dakota State University

Citation: More large, high-intensity forest fires likely in coming years (2017, April 6) retrieved 21 September 2024 from <https://phys.org/news/2017-04-large-high-intensity-forest-years.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.