

# Researchers find significant increase of invasive seaweed changing sea habitat

May 9 2017

---



A photo of what seaweed community looks like after introduction of invasive seaweed (*Dasysiphonia japonica*). Credit: Kristen Mello/UNH

Walking along the beaches of New England, it is easy to spot large

amounts of a fine red seaweed clogging the coastline, the result of sweeping changes in the marine environment occurring beneath the water. To further investigate, researchers at the University of New Hampshire looked at seaweed populations over the last 30 years in the Southwestern Gulf of Maine and found the once predominant and towering kelp seaweed beds are declining and more invasive, shrub-like species have taken their place, altering the look of the ocean floor and the base of the marine food chain.

In the study, recently published in the *Journal of Ecology*, researchers compared photos of sections of the sea floor, collected over 30 years, at several subtidal sites in the Southwestern Gulf of Maine. They also collected individual [seaweed species](#) to determine their complexity and the biodiversity of meso-invertebrates (smaller ocean species that fish and shellfish, such as crabs, feed on) associated with each [seaweed](#) species. The data showed that the seaweed community, as well as the number and types of small creatures, had significantly changed. The invasive fiber-like red seaweeds (*Dasysiphonia japonica*) had covered up to 90 percent of some areas, altering the visual landscape, and the newly created habitat structure now supported two to three times more small creatures at the base of the food chain.

"We were very surprised by what we saw," said Jennifer Dijkstra, research assistant professor in the Center for Coastal and Ocean Mapping at UNH and the lead author of the study. "In some areas, what was once a forest of tall blades of kelp with a high canopy height was now composed of bushy invasive seaweed species which had a much shorter canopy and a very different physical form."

Studies have found that kelp forests are one of the most productive systems in the ocean with high biodiversity and ecological function. They occur along the coastlines of most continents. Kelp provides a long three dimensional structure that offers protection and a source of food

for many juvenile [species](#) of fish (pollock, cod, and flounder), juvenile and adult shellfish (lobsters and crabs), seals and birds (terns and gulls).

"While the changing seascape has dramatically altered and increased the diversity and number of small creatures at the base of the marine food web, we still don't know how these changes in the ecosystem will propagate through the entire chain. Even though there may be more creatures at the base, it's not clear what their effects will be on fish or other crabs in the habitat, and how much protection the new landscape will provide," said Dijkstra.

Researchers say on-going studies are looking at the effects of the invasive types of seaweed and why they are so successful in the Gulf of Maine. They speculate that a number of events that relate to historical fishing practices, both commercial and recreational, combined with the warming waters in the Gulf of Maine may be increasing the negative effects of the growth of [kelp](#).

**More information:** Jennifer A. Dijkstra et al. Invasive seaweeds transform habitat structure and increase biodiversity of associated species, *Journal of Ecology* (2017). [DOI: 10.1111/1365-2745.12775](https://doi.org/10.1111/1365-2745.12775)

Provided by University of New Hampshire

Citation: Researchers find significant increase of invasive seaweed changing sea habitat (2017, May 9) retrieved 19 April 2024 from <https://phys.org/news/2017-05-significant-invasive-seaweed-sea-habitat.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.