


Precision control of superconductivity in atomic layers using magnetic molecules

May 15 2017



(a) Schematic diagram of a heterostructure comprising organic molecules and a superconducting atomic layer. (b)-(d) Scanning tunneling microscope images of samples. (b) Indium atomic layer (superconducting layer). (c) Phthalocyanine molecular layer grown on the indium atomic layer (manganese atom at the center of each molecule). (d) Phthalocyanine molecular layer grown on the indium atomic layer (copper atom at the center of each molecule). Credit: National Institute for Materials Science (NIMS)

A research team led by Shunsuke Yoshizawa, ICYS researcher, NIMS, Takashi Uchihashi, leader of the Surface Quantum Phase Materials Group, MANA, NIMS, Emi Minamitani, assistant professor, School of Engineering, University of Tokyo, Toshihiko Yokoyama, professor, IMS, NINS, and Kazuyuki Sakamoto, professor, Graduate School of Advanced Integration Science, Chiba University, succeeded in precisely controlling the transition temperature of atomic-scale-thick superconductors using magnetic organic molecules. The team also identified the control mechanism.

Atomic layer materials, including graphene, have been actively studied in recent years. In particular, much attention has been drawn to discoveries of superconducting atomic layer materials with a high transition temperature. These materials are superior to bulk materials in that their superconducting properties can be controlled through carrier doping of their surfaces/interfaces. However, it had been difficult to understand the mechanism of carrier doping at the microscopic level.

The research team recently succeeded for the first time in precisely controlling the transition temperature of superconducting [atomic layers](#) using [organic molecules](#). To achieve this, the team fabricated an ideal heterostructure consisting of a superconducting atomic layer and a layer of highly ordered organic molecules on top of the atomic layer. The creation of the heterostructure enabled the team to perform a detailed study on the mechanism behind the doping of atomic layer materials. Consequently, the analysis revealed that the metal atoms at the center of the organic molecules retained electron spins, which could generate magnetism. In addition, the team found that change in superconducting transition temperature is strongly influenced by competition between electron charge and spin in the organic molecules. Moreover, the team discovered that the spin effect is governed by the direction of electron orbitals, which can be viewed as "hidden degrees of freedom" in [molecules](#).

In light of these results, the researchers aim to greatly enhance [superconducting properties](#), i.e. superconducting [transition temperature](#), of these heterostructures . After such enhancement is made, the researchers hope to apply superconducting materials in a wide variety of fields in a manner so that the technology will help ease environmental/energy issues and support the sustainable development of society.

This study was published in *Nano Letters*, an American Chemical Society journal, on March 30, 2017.

More information: Shunsuke Yoshizawa et al. Controlled Modification of Superconductivity in Epitaxial Atomic Layer–Organic Molecule Heterostructures, *Nano Letters* (2017). [DOI: 10.1021/acs.nanolett.6b05010](#) Shunsuke Yoshizawa et al. Controlled Modification of Superconductivity in Epitaxial Atomic Layer–Organic Molecule Heterostructures, *Nano Letters* (2017). [DOI: 10.1021/acs.nanolett.6b05010](#)

Provided by National Institute for Materials Science

Citation: Precision control of superconductivity in atomic layers using magnetic molecules (2017, May 15) retrieved 19 September 2024 from <https://phys.org/news/2017-05-precision-superconductivity-atomic-layers-magnetic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.