


# Fractionating DNA—from hours to minutes

May 29 2017



Credit: University of Twente


Researchers of the University of Twente in The Netherlands developed a glass microchip for ultrafast separation and purification of DNA fragments. The chip, moreover, is easy to produce and cheap. The UT scientists published their results in *Microsystems and Nano Engineering*, a *Nature* publication.

The new chip is capable of fractionating DNA fragments within just a few minutes, while conventional approaches take hours. The chip does

this in high resolution and also purifies the fragments; it removes the other salts in the DNA sample. Tiny amounts of DNA, like in medical diagnostics or in forensics, will be sufficient. The fragments are typically in the size range of second generation DNA sequencing, the next step after the well-known Human Genome Project.

## Moving snake

UT scientist Burcu Gumuscu and her colleagues achieved this high speed and resolution by inventing a new approach to the common technique of gel electrophoresis. In the conventional approach, the DNA moves in a gel by an applied electric field. Larger fragments will move slower than small ones, and in this way size separation is possible. But the electric field has one direction and the molecules move in a [straight line](#). What about varying the field, the UT scientists thought. If you periodically apply an [electric field](#) of different magnitude in the perpendicular direction, the fragments will also respond to this. They don't just move in a straight line now, but somewhere in between the two field directions. As large fragments respond differently - like a snake moves - to the fields than small ones, fragments can be separated.



Credit: University of Twente

## **Cheap**

For collecting the fragments, the chip has arrays of micro channels on the sides of a square separation chamber of about 1 square centimeter. Next to that, it has a DNA reservoir and electrodes for applying the electric fields. The chip is relatively easy to produce using fundamental cleanroom techniques of the MESA+ NanoLab, and cheap. It is versatile as well, as the type and concentration of gel and the electric fields can be

adjusted to the application. Protein separation is one of the other possible application areas.

The research shows that DNA fragments between 0,5 kbp and 10 kpb (1 kbp= 1000 base pairs) can be fractionated within two minutes, at [high resolution](#). These are typically the smaller fragments, essential for the second generation DNA sequencing. Speeding up the separation is also a major step forward in medical applications.

The research has been done in the BIOS Lab-on-a-chip group, part of the MESA+ and MIRA research institutes of the University of Twente.

**More information:** Burcu Gumuscu et al. Exploiting biased reptation for continuous flow preparative DNA fractionation in a versatile microfluidic platform, *Microsystems & Nanoengineering* (2017). [DOI: 10.1038/micronano.2017.1](https://doi.org/10.1038/micronano.2017.1)

Provided by University of Twente

Citation: Fractionating DNA—from hours to minutes (2017, May 29) retrieved 19 September 2024 from <https://phys.org/news/2017-05-fractionating-dnafrom-hours-minutes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.