

'Harder, better, faster, stronger'-tethered soft exosuit reduces metabolic cost of running

May 31 2017



A system of actuation wires attached to the back of the exosuit provides assistive force to the hip joint during running. Credit: The Wyss Institute at Harvard University

What if running the 26.2 miles of a marathon only felt like running 24.9 miles, or if you could improve your average running pace from 9:14 minutes/mile to 8:49 minutes/mile without weeks of training?


Researchers at the Wyss Institute and John A. Paulson School of Engineering and Applied Sciences (SEAS) at Harvard University have demonstrated that a tethered soft exosuit can reduce the metabolic cost of running on a treadmill by 5.4% compared to not wearing the exosuit, bringing those dreams of high performance closer to reality. "Homo sapiens has evolved to become very good at distance running, but our results show that further improvements to this already extremely efficient system are possible," says corresponding author Philippe Malcolm, Ph.D., former Postdoctoral Research Fellow at the Wyss Institute and SEAS, and now Assistant Professor at the University of Nebraska, Omaha, where he continues to collaborate on this work. The study appears today in *Science Robotics*.

Running is a naturally more costly form of movement than walking, so any attempt to reduce its strain on the body must impose a minimal additional burden. The soft exosuit technology developed in the lab of Wyss Core Faculty member Conor Walsh, Ph.D., represents an ideal platform for assisted running, as its textile-based design is lightweight and moves with the body. A team of scientists in Walsh's lab led by Wyss Postdoctoral Fellow Giuk Lee, Ph.D. performed the study with an exosuit that incorporated flexible wires connecting apparel anchored to the back of the thigh and waist belt to an external actuation unit. As

subjects ran on a treadmill wearing the exosuit, the actuation unit pulled on the wires, which acted as a second pair of hip extensor muscles applying force to the legs with each stride. The metabolic cost was measured by analyzing the subjects' oxygen consumption and carbon dioxide production while running.

The team tested two different "assistance profiles," or patterns of wire-pulling: one based on human biology that applied force starting at the point of maximum hip extension observed in normal running, and one based on a simulation of exoskeleton-assisted running from [a group at Stanford University](#) that applied force slightly later in the running stride and suggested that the optimal point to provide assistive force might not be the same as the biological norm. Confirming this suspicion, Lee and colleagues found that the simulation-based profile outperformed the biology-based profile in reducing metabolic cost by about a factor of two. As the authors note in the paper, "Our finding supports a paradigm shift toward the concept that mimicking our current understanding of biology is not necessarily always optimal."

To clarify why applying force later in running stride improved metabolic expenditure so dramatically, the team analyzed what was happening to the subjects' other joints when their hip joints were being assisted by the actuating wires. They found that the simulation-based profile also affected knee extension and the forces between the foot and the ground, while the biology-based profile did not. "The biological profile only takes into account the amount of torque in the hip joint, but the human body is not a series of independently acting parts - it's full of muscles that act on multiple joints to coordinate movement," says Lee. "Applying force to the hip affects the whole body system, and we need to consider that in order to give the best assistance." While the study's results suggest that the increased biomechanical involvement of the simulation-based assistance profile is responsible for the reduction in metabolic cost, further biomechanical analyses are needed to confirm its findings.



A system of actuation wires attached to the back of the exosuit provides assistive force to the hip joint during running. Credit: The Wyss Institute at Harvard University

The team hopes to continue this research to reduce the metabolic cost of running even more. "We only tested two actuation profiles in this study, so it will be interesting to see how much more the cost of running can be reduced with further optimization of the system," says Malcolm. "Our goal is to develop a portable system with a high power-to-weight ratio so that the benefit of using the suit greatly offsets the cost of wearing it. We believe this technology could augment the performance of recreational athletes and/or help with recovery after injury," adds Lee. The days of a battery-powered exosuit for high-performance runners are still beyond the horizon, as the actuator unit (including motors, electronics, and power supply) in this study was off-board, but the

authors say technology is moving toward making an untethered assistive [exosuit](#) possible in the near future. "This study is a great example of how our team can quickly leverage its experience and expertise in wearable technology and biomechanics to demonstrate how human performance can be improved in new application areas," says Walsh.

"Our work at the Wyss is biologically inspired and, as shown by Conor's team's advance, sometimes we learn that a conventional biological view of a mechanism is not really how living systems work; we need to place individual components - in this case, a human hip joint - in context of the living whole. When we do this right, transformative new technologies emerge," says the Wyss' Founding Director Donald Ingber, M.D., Ph.D., the Judah Folkman Professor of Vascular Biology at Harvard Medical School (HMS) and the Vascular Biology Program at Boston Children's Hospital, as well as Professor of Bioengineering at Harvard's School of Engineering and Applied Sciences (SEAS). "Studies like this are also a great example of the power of thinking beyond the limitations that Nature has handed us and asking, 'Can we do better?'"

More information: G. Lee et al., "Reducing the metabolic cost of running with a tethered soft exosuit," *Science Robotics* (2017).
robotics.sciencemag.org/lookup.../scirobotics.aan6708

Provided by Harvard University

Citation: 'Harder, better, faster, stronger'-tethered soft exosuit reduces metabolic cost of running (2017, May 31) retrieved 26 April 2024 from <https://phys.org/news/2017-05-harder-faster-stronger-tethered-soft-exosuit.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
