


A possible explanation for why no intermediate sized black holes have been found

27 June 2017, by Bob Yirka



This computer-simulated image shows a supermassive black hole at the core of a galaxy. The black region in the center represents the black hole's event horizon, where no light can escape the massive object's gravitational grip. The black hole's powerful gravity distorts space around it like a funhouse mirror. Light from background stars is stretched and smeared as the stars skim by the black hole. Credit: NASA, ESA, and D. Coe, J. Anderson, and R. van der Marel (STScI)

(Phys.org)—A pair of researchers, one with the Weizmann Institute of Science in Israel the other with Princeton University in the U.S. has come up with a possible explanation for the inability of space scientists to find any intermediate-sized black holes. In their paper published in the journal *Nature Astronomy*, Tal Alexander and Ben Bar-Or offer an outline of current theories regarding how black holes in general are believed to develop,

theories about the early universe, and finally, their ideas on how current theories may lead to an explanation for the dearth of intermediate sized black holes.

Despite their popularity both in real science and [science](#) fiction, the concept of a black hole has only been around for a hundred years, as predicted by Albert Einstein. The term itself did not come into use until 1967, and it was just 46 years ago that the first one was identified. Physicists have learned much about [black holes](#), though, particularly over the past several decades, but one glaring mystery remains—why have scientists never been able to spot one in the intermediate size range? Many large and small ones have been found, but none in the middle. In their paper, Alexander and Bar-Or suggest that our inability to find mid-sized black holes does not mean they do not exist—we just might have to look for them in other places.

The researchers suggest that there may be fewer, or even no intermediate black holes today because of the way they grow—prior research has indicated that they likely get bigger as they "eat" stars. If so, they contend, then their calculations show black holes growing at a rate of one solar mass per 10,000 years. That means, they note, that if a black hole was born soon after the universe came into existence approximately 13.8 billion years ago (which is believed to be the case for large black holes), a black hole would have grown well past the intermediate stage by now. They also suggest that if intermediate-sized black holes do exist, they might reside in dense parts of the sky, which would make them very hard to spot. But such problems might be overcome soon, they note, as new technology has begun detecting gravitational waves.

More information: Tal Alexander et al. A

universal minimal mass scale for present-day central black holes, *Nature Astronomy* (2017). DOI: [10.1038/s41550-017-0147](https://doi.org/10.1038/s41550-017-0147)

Abstract

The early stages of massive black hole growth are poorly understood. High-luminosity active galactic nuclei at very high redshift z further imply rapid growth soon after the Big Bang. Suggested formation mechanisms typically rely on the extreme conditions found in the early Universe (very low metallicity, very high gas or star density). It is therefore plausible that these black hole seeds were formed in dense environments, at least a Hubble time ago ($z > 1.8$ for a look-back time of $t_H = 10$ Gyr). Intermediate-mass black holes (IMBHs) of mass $M \sim 10^2 - 10^5$ solar masses, M , are the long-sought missing link between stellar black holes, born of supernovae, and massive black holes, tied to galaxy evolution by empirical scaling relations. The relation between black hole mass, M , and stellar velocity dispersion, σ , that is observed in the local Universe over more than about three decades in massive black hole mass, correlates M and σ on scales that are well outside the massive black hole's radius of dynamical influence, $r \sim GM/\sigma^2 \sim GM/\sigma^2$. We show that low-mass black hole seeds that accrete stars from locally dense environments in galaxies following a universal M/σ relation grow over the age of the Universe to be above $M \gtrsim 3 \times 10^5 M_\odot$ (5% lower limit), independent of the unknown seed masses and formation processes. The mass M_0 depends weakly on the uncertain formation redshift, and sets a universal minimal mass scale for present-day black holes. This can explain why no IMBHs have yet been found, and it implies that present-day galaxies with σ

APA citation: A possible explanation for why no intermediate sized black holes have been found (2017, June 27) retrieved 21 September 2021 from <https://phys.org/news/2017-06-explanation-intermediate-sized-black-holes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.