

# Extreme low-oxygen eddies in the Atlantic produce greenhouse gases

July 7 2017

---



Credit: CC0 Public Domain

Oxygen in the seawater is not only vital to most marine organisms, its concentrations also affect the chemistry of the ocean and that of the atmosphere above. In oceanic regions with very little oxygen, for

example, large amounts of the potent greenhouse gas nitrous oxide, also called laughing gas, are produced via biogeochemical processes and can then be released to the atmosphere.

Even though a natural moderate [oxygen](#) minimum zone (OMZ) exists along some of the eastern boundaries of the Atlantic Ocean, the Atlantic OMZ, unlike the OMZs of the Indian and Pacific oceans, was not considered to be a region of extremely low oxygen concentrations. New findings by an international research team led by the Kiel Excellence Cluster "Future Ocean" and the GEOMAR Helmholtz Center for Ocean Research Kiel, however, now imply that this picture has to be corrected. This study was published yesterday in the Nature Publishing Group journal *Scientific Reports*.

"The reason why the extremely low-oxygen regions in the Atlantic have so far escaped research is simple: they are relatively small and mobile in contrast to the well-known, large and stationary oxygen minimum zones", explains Dr. Damian Grundle from the Bermuda Institute of Ocean Sciences, first author of the current study and, until recently, a scientist at the "Future Ocean"/GEOMAR. Extreme low oxygen concentrations occur in the Atlantic in [ocean](#) eddies of up to 100 kilometers in diameter, which migrate westward across the ocean from the West African coast. Such eddies are difficult to detect with conventional observation methods, and require a combination of satellite, glider and ship-based observations.

The first observation of a low oxygen eddy in the Atlantic Ocean, however, was detected by the Cape Verde Ocean Observatory, a project which includes an ocean observation mooring north of the Cape Verdean island of São Vicente. "We had a first indication of the existence of these special eddies but still no exact information from their interior," reports the marine chemist Dr. Björn Fiedler from GEOMAR, who led the project.

With the financial support of the Kiel Cluster of Excellence "The Future Ocean", an interdisciplinary group of scientists waited for their next chance. It came in 2014: by means of satellite observation they discovered a potential low oxygen eddy, which formed off the coast of Mauritania and migrated towards Cape Verde. From there, the team deployed autonomous sensor carriers, so-called gliders, to intercept the eddy. As it approached the islands, the researchers were also able to take water samples directly from the eddy's center with the Cape Verdean research vessel ISLANDIA.

"Chance also helped us. At the time we investigated the eddy the German research vessel METEOR conducted a long planned expedition for the Collaborative Research Centre 754 Kiel off Cape Verde. We quickly convinced our colleagues to sample the eddy, too," says Dr. Fiedler. He adds: "Without the good infrastructure on the Cape Verde Islands and the long-term cooperation with our colleagues there, this campaign would not have been possible."

Afterwards the obtained data and water samples were evaluated physically, biogeochemically and biologically. "In a whole series of publications, we were able to gain exciting new insights into the hitherto unknown phenomenon in the Atlantic," says Dr. Fiedler.

The latest study, now published in *Scientific Reports*, demonstrates that at the core of the eddy, the highest levels of the greenhouse gas nitrous oxide ever measured in the open Atlantic were found in only 100 meters of water depth. This can be attributed to processes that can deplete the plant nutrient nitrogen from the ocean, producing, among other things, nitrous oxide in large quantities, and further points out that scientists should now consider revising our understanding of the chemical cycles in the Atlantic.

**More information:** D. S. Grundle et al, Low oxygen eddies in the

eastern tropical North Atlantic: Implications for N<sub>2</sub>O cycling, *Scientific Reports* (2017). [DOI: 10.1038/s41598-017-04745-y](https://doi.org/10.1038/s41598-017-04745-y)

Provided by Helmholtz Association of German Research Centres

Citation: Extreme low-oxygen eddies in the Atlantic produce greenhouse gases (2017, July 7)  
retrieved 20 September 2024 from  
<https://phys.org/news/2017-07-extreme-low-oxygen-eddies-atlantic-greenhouse.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.