


Magnetic fields in massive star formation cores

August 7 2017



A far-infrared image of the long filament of star formation activity known as

DR21, seen here in emission by the Herschel Space Telescope. A study of the magnetic field along the filament and around six star-forming cores within it finds that magnetic effects are primarily important during the early stages of star formation. Credit: ESA-Herschel

Studies of molecular clouds have revealed that star formation usually occurs in a two-step process. First, supersonic flows compress the clouds into dense filaments light-years long, after which gravity collapses the densest material in the filament into cores. In this scenario, massive cores (each more than about 20 solar masses) preferentially form at intersections where filaments cross, producing sites of clustered star formation. The process sounds reasonable and is expected to be efficient, but the observed rate of star formation in dense gas is only a few percent of the rate expected if the material really were freely collapsing. To solve the problem, astronomers have proposed that magnetic fields support the cores against the collapse induced by self-gravity.

Magnetic fields are difficult to measure and difficult to interpret. CfA astronomers Tao-Chung Ching, Qizhou Zhang, and Josep Girat led a team that used the Submillimeter Array to study six dense cores in a nearby [star formation](#) region in Cygnus. They measured the field strengths from the polarization of the millimeter radiation; elongated dust grains are known to be aligned by magnetic fields and to scatter light with a preferred polarization direction. The scientists then correlated the field direction in these cores with the field direction along the filament out of which the cores developed.

The astronomers find that the [magnetic field](#) along the filament is well-ordered and parallel to the structure, but at the cores themselves the field direction is much more complex, sometimes parallel and sometimes

perpendicular. They conclude that during the formation of the cores the magnetic fields, at least at small scales, become unimportant compared to turbulence and infall. Although the field may play an important role as the filament initially collapses, once the dense cores develop the local kinematics from infall and gravitational effects become more important.

More information: Tao-Chung Ching et al. Magnetic Fields in the Massive Dense Cores of the DR21 Filament: Weakly Magnetized Cores in a Strongly Magnetized Filament, *The Astrophysical Journal* (2017).

[DOI: 10.3847/1538-4357/aa65cc](https://doi.org/10.3847/1538-4357/aa65cc)

Provided by Harvard-Smithsonian Center for Astrophysics

Citation: Magnetic fields in massive star formation cores (2017, August 7) retrieved 20 September 2024 from

<https://phys.org/news/2017-08-magnetic-fields-massive-star-formation.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.