


Researchers identify transport mechanism for synaptic junction reinforcement

7 August 2017



Fluorescent in situ hybridisation in hippocampal neurons to visualize intron containing Calmodulin3 mRNA localization (in White). The dendritic processes of the neurons are colored green while their nuclei are blue. Credit: T. Sharangdhar and M. Kiebler, LMU

Learning involves constant restructuring of nerve-cell connections. This requires specialized transport systems to ensure that these specific synapses can structurally and functionally be modified. One such delivery mechanism has now been characterized.

At the cellular level, learning and memory require continuous readjustment of the strength of synaptic junctions – the functional connections between nerve cells. These modifications are achieved by the incorporation into individual [synapses](#) of proteins that modulate the efficiency of intercellular signal transmission. The blueprints for the synthesis of these molecules, in the form of messenger RNAs (mRNAs), are delivered to active synapses by a dedicated transport system. An international team led by LMU biochemist Professor Michael Kiebler has now demonstrated that a key factor involved in this transport binds specifically to regions of its mRNA cargo that lack protein-coding information, and that release of the

RNAs to enable protein synthesis is dependent on [synaptic activity](#). The new findings appear online in the journal *EMBO Reports*.

Nerve [cells](#) typically consist of a cell body (in which the nucleus resides), a long fiber (the axon) which conducts outgoing nerve impulses, and a set of shorter projections (dendrites) near the cell body that serve as antennas for incoming signals. The instructions for the synthesis of proteins are stored in the DNA in the [cell nucleus](#), are first 'transcribed' into mRNAs, and exported from the nucleus to the ribosomes in the cell cytoplasm. Ribosomes are the molecular machines responsible for the translation of mRNA sequences into the amino-acid sequences of the specified newly synthesized protein. In [nerve cells](#), ribosomes are also found close to the synapses of the dendrites, thus enabling proteins that are required for synaptic function to be produced close to their sites of action. This localized delivery system also enables particular synapses to be specifically modified without concomitant alteration of their neighbors.

Kiebler and his group had previously shown that the brain-specific RNA-binding protein Stau2 (Stau2) plays a central role in the transport of mRNAs from the cell nucleus to synaptic ribosomes. "However, it remained unclear how Stau2 recognizes the mRNAs it conveys to the synapses, and how these are translated into proteins that modify specific synapses in an activity-dependent manner," Kiebler explains.

In the new study, the researchers show that Stau2 binds specifically to a so-called intron sequence found near the end of a specific subset of mRNAs – in a region that contains no coding information, but serves a regulatory function instead. Introns are regions in the so-called primary transcript transcribed from the nuclear DNA, which are either removed (by a process called alternative splicing) or selectively retained prior to export of the processed mRNA. "Most introns found in mRNAs

are located in protein-coding regions and their selective retention permits different versions – so-called isoforms – of a protein to be synthesized from the same precursor RNA. It is however quite unusual to find an intron retained in the untranslated region of an mRNA," says Tejaswini Sharangdhar, first author of the study. Moreover, Stau2 fails to bind to the mRNA isoform that lacks the intron. Consequently, this RNA isoform is not transported to the dendrites for translation. Thus the intron in the non-coding region serves as one of the regulatory markers which ensure that mRNA is recognized by Stau2 and transported to synapses.

"Our second surprising discovery is that this transport process is dependent on synaptic activity," says Kiebler: The intron-containing mRNA is transported from one synapse to the next – but will only be released for translation in the vicinity of synapses that are momentarily active. Hence, synaptic activity stimulates the docking of the mRNA to specific synapses, and its protein product marks "learning" synapses. Here too, Stau2 plays an essential role, for in the absence of the protein, none of the mRNA reaches the dendrites.

More information: Tejaswini Sharangdhar et al. A retained intron in the 3'UTR of Calm3 mRNA mediates its Stau2 and activity-dependent localization to neuronal dendrites, *EMBO reports* (2017). DOI: [10.15252/embr.201744334](https://doi.org/10.15252/embr.201744334)

Provided by Ludwig Maximilian University of Munich

APA citation: Researchers identify transport mechanism for synaptic junction reinforcement (2017, August 7) retrieved 16 June 2019 from <https://phys.org/news/2017-08-mechanism-synaptic-junction.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.