

# Meet the hominin species that gave us genital herpes

2 October 2017



Cast of a *P. boisei* skull, used for teaching at Cambridge University Credit: Louise Walsh

Two herpes simplex viruses infect primates from unknown evolutionary depths. In modern humans these viruses manifest as cold sores (HSV1) and genital herpes (HSV2).

Unlike HSV1, however, the earliest proto-humans did not take HSV2 with them when our ancient lineage split from chimpanzee precursors around 7 million years ago. Humanity dodged the [genital herpes](#) bullet - almost.

Somewhere between 3 and 1.4 million years ago, HSV2 jumped the [species](#) barrier from African apes back into human ancestors - probably through an intermediate hominin species unrelated to humans. Hominin is the zoological 'tribe' to which our species belongs.

Now, a team of scientists from Cambridge and Oxford Brookes universities believe they may have identified the culprit: *Paranthropus boisei*, a heavyset bipedal hominin with a smallish brain and dish-like face.

In a study published today in the journal *Virus Evolution*, they suggest that *P. boisei* most likely contracted HSV2 through scavenging ancestral chimp meat where savannah met forest - the infection seeping in via bites or open sores.

Hominins with HSV1 may have been initially protected from HSV2, which also occupied the mouth. That is until HSV2 "adapted to a different mucosal niche" say the scientists. A niche located in the genitals.

Close contact between *P. boisei* and our ancestor *Homo erectus* would have been fairly common around sources of water, such as Kenya's Lake Turkana. This provided the opportunity for HSV2 to boomerang into our bloodline.

The appearance of *Homo erectus* around 2 million years ago was accompanied by evidence of hunting and butchery. Once again, consuming "infected material" would have transmitted the virus - only this time it was *P. boisei* being devoured.

"Herpes infect everything from humans to coral, with each species having its own specific set of viruses," said senior author Dr Charlotte Houldcroft, a virologist from Cambridge's Department of Archaeology.

"For these viruses to jump species barriers they need a lucky genetic mutation combined with significant fluid exchange. In the case of early hominins, this means through consumption or intercourse - or possibly both."

"By modelling the available data, from fossil records to viral genetics, we believe that *Paranthropus boisei* was the species in the right place at the right time to both contract HSV2 from ancestral chimpanzees, and transmit it to our earliest ancestors, probably *Homo erectus*."

When researchers from University of California,

San Diego, published findings suggesting HSV2 had jumped between hominin species, Houldcroft became curious.

While discussing genital herpes over dinner at Kings College, Cambridge, with fellow academic Dr Krishna Kumar, an idea formed. Kumar, an engineer who uses Bayesian network modelling to predict city-scale infrastructure requirements, suggested applying his techniques to the question of ancient HSV2.

Houldcroft and her collaborator Dr Simon Underdown, a human evolution researcher from Oxford Brookes, collated data ranging from fossil finds to herpes DNA and ancient African climates. Using Kumar's model, the team generated HSV2 transmission probabilities for the mosaic of hominin species that roamed Africa during "deep time".

"Climate fluctuations over millennia caused forests and lakes to expand and contract," said Underdown. "Layering climate data with fossil locations helped us determine the species most likely to come into contact with ancestral chimpanzees in the forests, as well as other hominins at water sources."

Some promising leads turned out to be dead ends. *Australopithecus afarensis* had the highest probability of proximity to ancestral chimps, but geography also ruled it out of transmitting to human ancestors.

Ultimately, the researchers discovered the key player in all the scenarios with higher probabilities to be *Parathropus boisei*. A genetic fit virally who was found in the right places to be the herpes intermediary, with *Homo erectus* - and eventually us - the unfortunate recipients.

"Once HSV2 gains entry to a species it stays, easily transferred from mother to baby, as well as through blood, saliva and sex," said Houldcroft.

"HSV2 is ideally suited to low density populations. The genital herpes virus would have crept across Africa the way it creeps down nerve endings in our sex organs - slowly but surely."

The team believe their methodology can be used to unravel the transmission mysteries of other ancient diseases - such as human pubic lice, also introduced via an intermediate [hominin](#) from ancestral gorillas over 3 million years ago.

**More information:** Simon J. Underdown et al, Network analysis of the hominin origin of Herpes Simplex virus 2 from fossil data, *Virus Evolution* (2017). [DOI: 10.1093/ve/vex026](https://doi.org/10.1093/ve/vex026)

Provided by University of Cambridge

APA citation: Meet the hominin species that gave us genital herpes (2017, October 2) retrieved 27 November 2022 from <https://phys.org/news/2017-10-hominin-species-gave-genital-herpes.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*