


Physicists open the door to the first direct measurement of Berry curvature in solid matter

November 22 2017, by Julie Cohen



This artist's rendition features Berry curvature represented by the twisting

ribbons at the top. Credit: Brian Long

Berry curvature may not be the most well-known scientific concept, but to many physicists, its direct measurement is something akin to a holy grail.

A powerful unifying principle in several branches of classical and quantum physics, Berry curvature is a strange and elusive quantum mechanical property of solids. It governs the dynamics of the motion of charges in semiconductors yet itself cannot be directly measured.

If it could be, the resulting calculation could lead to new materials for quantum computing.

Now, UC Santa Barbara physicists have opened the door to the first direct measurement of Berry curvature in solid matter. Their work, published in the journal *Physical Review X*, builds on a previous UCSB paper in which they describe experiments resulting in an electron-hole recollision achieved by aiming high- and low-frequency laser beams at a [semiconductor](#) made of gallium arsenide.

For the new paper, scientists from UCSB's Sherwin Group collaborated with colleagues in China, at Princeton University and at the U.S. Naval Research Laboratory to improve on the previous experiment. They discovered a surprising new phenomenon using the same semiconductor subjected to extremely strong laser fields oscillating nearly 1 trillion times per second (1 terahertz). Called dynamical birefringence, this phenomenon can be used to investigate Berry curvature.

"When we originally did the experiment, we could only detect one sideband at a time and the samples were very fragile and more difficult

to work with," explained corresponding author Mark Sherwin, director of UCSB's Institute for Terahertz Science and Technology and a professor in the Department of Physics.

Hunter Banks, lead author of the new paper, installed a camera that allowed the team to see all the sidebands simultaneously, which decreased the experiment's duration and increased its sensitivity. He also improved how the samples were mounted and increased the strength of the terahertz electric field that could be applied.

These enhancements revealed sidebands separated by as much as 90 times the photon energy of the terahertz—more than three times the amount in the original experiment. Sherwin noted that an increased quantity of sidebands allowed the team to learn more about the semiconductor. "As far as we know, this huge number of sidebands is the highest-order nonlinear optical process in solids," he said.

Generated by a unique laser housed in a dedicated building at UCSB, these experiments drive thin layers of semiconductor while they are illuminated by weak infrared light. The infrared light is polarized in one of two ways: either parallel or perpendicular to the terahertz field.

"The [infrared light](#) transmitted through the semiconductor exhibits a rainbowlike spectrum containing dozens of frequencies, or sidebands," Sherwin explained. "Unexpectedly, the sidebands are usually stronger when the infrared beam is polarized perpendicular to the terahertz field. Somehow the [terahertz](#) is actually defining an axis that acts like a polarizing one. We call this phenomenon dynamical birefringence, and it arises as a direct consequence of Berry curvature."

It also creates possibilities for applications in new classes of electronic and optical devices.

"We're planning to turn dynamical birefringence into a direct measurement of Berry curvature," Sherwin explained. "Once you can measure something that's a basic property of a solid, then, when you're designing new materials, you can optimize the Berry curvature for a particular device."

To navigate through an annotated 3-D rendering with virtual reality capability, [click here](#).

More information: Hunter B. Banks et al, Dynamical Birefringence: Electron-Hole Recollisions as Probes of Berry Curvature, *Physical Review X* (2017). [DOI: 10.1103/PhysRevX.7.041042](https://doi.org/10.1103/PhysRevX.7.041042)

Provided by University of California - Santa Barbara

Citation: Physicists open the door to the first direct measurement of Berry curvature in solid matter (2017, November 22) retrieved 20 September 2024 from <https://phys.org/news/2017-11-physicists-door-berry-curvature-solid.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--