


ALMA finds massive primordial galaxies swimming in vast ocean of dark matter

6 December 2017



A composite image showing ALMA data (red) of the two galaxies of SPT0311-58. These galaxies are shown over a background from the Hubble Space Telescope (blue and green). The ALMA data show the two galaxies' dusty glow. The image of the galaxy on the right is distorted by gravitational lensing. The nearer foreground lensing galaxy is the green object between the two galaxies imaged by ALMA. Credit: ALMA (ESO/NAOJ/NRAO), Marrone, et al.; B. Saxton (NRAO/AUI/NSF); NASA/ESA Hubble

Astronomers expect that the first galaxies, those that formed just a few hundred million years after the Big Bang, would share many similarities with some of the dwarf galaxies we see in the nearby universe today. These early agglomerations of a few billion stars would then become the building blocks of the larger galaxies that came to dominate the universe after the first few billion years.

Ongoing observations with the Atacama Large

Millimeter/submillimeter Array (ALMA), however, have discovered surprising examples of massive, star-filled galaxies seen when the cosmos was less than a billion years old. This suggests that smaller galactic building blocks were able to assemble into large galaxies quite quickly.

The latest ALMA observations push back this epoch of massive-galaxy formation even further by identifying two [giant galaxies](#) seen when the universe was only 780 million years old, or about 5 percent its current age. ALMA also revealed that these uncommonly large galaxies are nestled inside an even-more-massive cosmic structure, a halo of [dark matter](#) several trillion times more massive than the sun.

The two galaxies are in such close proximity—less than the distance from the Earth to the center of our galaxy—that they will shortly merge to form the largest galaxy ever observed at that period in cosmic history. This discovery provides new details about the emergence of large galaxies and the role that dark [matter](#) plays in assembling the most massive structures in the universe.


The researchers report their findings in the journal *Nature*.

"With these exquisite ALMA observations, astronomers are seeing the most massive galaxy known in the first billion years of the universe in the process of assembling itself," said Dan Marrone, associate professor of astronomy at the University of Arizona in Tucson and lead author on the paper.

Astronomers are seeing these galaxies during a period of cosmic history known as the Epoch of Reionization, when most of intergalactic space was suffused with an obscuring fog of cold hydrogen gas. As more stars and galaxies formed, their energy eventually ionized the hydrogen between the galaxies, revealing the universe as we see it today.

"We usually view that as the time of little galaxies working hard to chew away at the neutral intergalactic medium," said Marrone. "Mounting observational evidence with ALMA, however, has helped to reshape that story and continues to push back the time at which truly massive galaxies first emerged in the universe."

The galaxies that Marrone and his team studied, collectively known as SPT0311-58, were originally identified as a single source by the South Pole Telescope. These first observations indicated that this object was very distant and glowing brightly in infrared light, meaning that it was extremely dusty and likely going through a burst of star formation. Subsequent observations with ALMA revealed the distance and dual nature of the object, clearly resolving the pair of interacting galaxies.



Artist impression of a pair of galaxies from the very early universe. Credit: NRAO/AUI/NSF; D. Berry

To make this observation, ALMA had some help from a gravitational lens, which provided an observing boost to the telescope. Gravitational lenses form when an intervening massive object, like a galaxy or galaxy cluster, bends the light from more distant galaxies. They do, however, distort the appearance of the object being studied, requiring sophisticated computer models to reconstruct the image as it would appear in its unaltered state.

This "de-lensing" process provided intriguing details

about the galaxies, showing that the larger of the two is forming stars at a rate of 2,900 solar masses per year. It also contains about 270 billion times the mass of our sun in gas and nearly 3 billion times the mass of our sun in dust. "That's a whopping large quantity of dust, considering the young age of the system," noted Justin Spilker, a recent graduate of the University of Arizona and now a postdoctoral fellow at the University of Texas at Austin.

The astronomers determined that this galaxy's rapid star formation was likely triggered by a close encounter with its slightly smaller companion, which already hosts about 35 billion solar masses of stars and is increasing its rate of starburst at the breakneck pace of 540 solar masses per year.

The researchers note that galaxies of this era are "messier" than the ones we see in the nearby universe. Their more jumbled shapes would be due to the vast stores of gas raining down on them and their ongoing interactions and mergers with their neighbors.

The new observations also allowed the researchers to infer the presence of a truly massive [dark matter halo](#) surrounding both galaxies. Dark matter provides the pull of gravity that causes the universe to collapse into structures (galaxies, groups and clusters of galaxies, etc.).

"If you want to see if a galaxy makes sense in our current understanding of cosmology, you want to look at the dark matter halo—the collapsed dark matter structure—in which it resides," said Chris Hayward, associate research scientist at the Center for Computational Astrophysics at the Flatiron Institute in New York City. "Fortunately, we know very well the ratio between dark matter and normal matter in the [universe](#), so we can estimate what the dark matter halo mass must be."

By comparing their calculations with current cosmological predictions, the researchers found that this halo is one of the most massive that should exist at that time.

"There are more galaxies discovered with the South Pole Telescope that we're following up on," said Joaquin Vieira of the University of Illinois at Urbana-

Champaign, "and there is a lot more survey data that we are just starting to analyze. Our hope is to find more objects like this, possibly even more distant ones, to better understand this population of extreme dusty galaxies and especially their relation to the bulk population of galaxies at this epoch."

"In any case, our next round of ALMA observations should help us understand how quickly these [galaxies](#) came together and improve our understanding of massive galaxy formation during reionization," added Marrone.

More information: D. P. Marrone et al, Galaxy growth in a massive halo in the first billion years of cosmic history, *Nature* (2017). [DOI: 10.1038/nature24629](#)

Provided by National Radio Astronomy Observatory

APA citation: ALMA finds massive primordial galaxies swimming in vast ocean of dark matter (2017, December 6) retrieved 25 September 2021 from <https://phys.org/news/2017-12-alma-massive-primordial-galaxies-vast.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.