


New catalyst for making fuels from shale gas

8 January 2018



STM imaging of reaction intermediates on Cu(111) and Pt/Cu SAA surfaces. Credit: Sykes

Methane in shale gas can be turned into hydrocarbon fuels using an innovative platinum and copper alloy catalyst, according to new research led by UCL (University College London) and Tufts University.

Platinum or nickel are known to break the [carbon-hydrogen bonds](#) in methane found in [shale gas](#) to make [hydrocarbon fuels](#) and other useful chemicals. However, this process causes 'coking' - the metal becomes coated with a carbon layer rendering it ineffective by blocking reactions from happening at the surface.

The new alloy [catalyst](#) is resistant to coking, so it retains its activity and requires less energy to break the bonds than other materials.

Currently, methane reforming processes are extremely energy intense, requiring temperatures of about 900 degrees Celsius. This new material could lower this to 400 degrees Celsius, saving

energy.

The study, published today in *Nature Chemistry*, demonstrates the benefits of the new highly diluted alloy of [platinum](#) in copper - a single atom alloy - in making useful chemicals from small hydrocarbons.

A combination of [surface science](#) and catalysis experiments and powerful computing techniques were used to investigate the performance of the alloy. These showed that the platinum breaks the carbon-hydrogen bonds, and the copper helps couple hydrocarbon molecules of different sizes, paving the way towards conversion to fuels.

Study co-lead author, Professor Michail Stamatakis (UCL Chemical Engineering), said: "We used supercomputers to model how the reaction happens - the breaking and making of bonds in small molecules on the catalytic alloy surface, and also to predict its performance at large scales. For this, we needed access to hundreds of processors to simulate thousands of reaction events."

While UCL researchers traced the reaction using computers, Tufts chemists and chemical engineers ran surface science and micro-reactor experiments to demonstrate the viability of the new catalyst - atoms of platinum dispersed in a copper surface - in a practical setting. They found the [single atom alloy](#) was very stable and only required a tiny amount of platinum to work.

Study lead, Professor Charles Sykes of the Department of Chemistry in Tufts University's School of Arts & Sciences, said: "Seeing is believing, and our scanning tunnelling microscope allowed us to visualise how single platinum atoms were arranged in copper. Given that platinum is over \$1,000 an ounce, versus copper at 15 cents, a significant cost saving can be made."

Together, the team shows that less energy is needed for the alloy to help break the bonds

between carbon and hydrogen atoms in methane and butane, and that the alloy is resistant to coking, opening up new applications for the material.

Study co-lead author, Distinguished Professor Maria Flytzani-Stephanopoulos of the Department of Chemical and Biological Engineering in Tufts University's School of Engineering, said: "While model catalysts in surface science experiments are essential to follow the structure and reactivity at the atomic scale, it is exciting to extend this knowledge to realistic nanoparticle catalysts of similar compositions and test them under practical conditions, aimed at developing the catalyst for the next step - industrial application."

The team now plan on developing further catalysts that are similarly resistant to the coking that plagues metals traditionally used in this and other chemical processes.

More information: Pt/Cu single-atom alloys as coke-resistant catalysts for efficient C–H activation, *Nature Chemistry* (2018). DOI: 10.1038/nchem.2915 , [nature.com/articles/doi:10.1038_nchem.2915](https://doi.org/10.1038/nchem.2915)

Provided by University College London

APA citation: New catalyst for making fuels from shale gas (2018, January 8) retrieved 21 June 2021 from <https://phys.org/news/2018-01-catalyst-fuels-shale-gas.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.