

Will the weather get worse in 2018? What the experts say

January 12 2018, by Lindsay Beevers



Credit: EPA, CC BY-SA

Last year unleashed some catastrophic weather across the world. At the beginning of 2017, Australia experienced one of the [hottest summers on record](#) in Sydney and Brisbane, followed by a killer [summer heatwave](#) across southern Europe and [wildfires](#) triggered by heat in California.

The Atlantic hurricane season was particularly active, recording three mighty category 5 hurricanes – [Harvey](#), [Irma](#) and [Maria](#) – resulting in

significant casualties. The cost of the damage across the American continent was in excess of [\\$260 billion](#).

At the same time, the 2017 monsoon season brought considerable rains to the Indian subcontinent, and resulted in [devastating floods](#) in parts of India, Pakistan, Nepal and Bangladesh (one of the most flood vulnerable countries in the world), causing more than 1,000 deaths.

At the end of 2017, we could look back at the statistics and see that globally it was the warmest [non-El Nino](#) year on record. El-Nino is the warming part of the [El-Nino Southern Oscillation](#) (ENSO) cycle which affects global ocean and atmospheric currents and can create powerful storms and hurricanes.

In a world that is getting warmer, what does that mean and what might 2018 hold for [weather](#) across the globe? Are we destined to see more and more of these "unusual" and catastrophic weather events which result in significant devastation? What effect will [climate change](#) have on our weather and will this become the new norm?



Monsoon flooding in the Himalayan foothills of India, Bangladesh and Nepal affected more than 40m people last year. Credit: EPA, CC BY-SA

Looking at the facts

First, we need to untangle the [difference](#) between weather and [climate](#) change, something president Trump seems [to confuse](#). In a nutshell, the difference is time. Weather is the conditions in the atmosphere over a short period of time. Climate is how the atmosphere behaves over a longer period of time. When we talk about climate change, that generally means changes in long-term averages of daily levels of temperature and rainfall. So we may see a change in average or typical weather over a number of years, but we can still experience extremes in any one year.

Recently, scientists have used [robust event attribution](#) where natural or human influences on particular events are studied to understand the role [climate change](#) may play in particular weather events. This can help support future regional contingency planning.

This also helps us to understand the role climate change has played in past events such as heatwaves, wildfires, droughts, or extreme flooding and some of their large-scale drivers such as ocean and atmospheric patterns of warming and cooling, like El-Nino.

Looking at the past there are many factors which control the onset, spread and eventual impact of a big weather event. But not all of these factors are climatic and many relate to things such as urbanisation, engineering interventions or land-use changes.

What the future looks like

According to the Intergovernmental Panel on Climate Change [IPCC](#), an international body set up to assess the science of climate change, we can continue to expect [an increase](#) in the average global temperature. That means we will be experiencing warmer years in the future.

But at the same time, we may see changes to the extremes, which could become more frequent in the case of high temperature or heavy rainfall, or less frequent in the case of extreme cold. This means that the distribution, occurrence and expected averages of our weather (for example, temperature and rain) throughout the year may change, resulting in warmer years on average with more extreme hot days, and fewer extreme cold days in the future.

This pattern has a direct link to such phenomena as heatwaves, which are caused by more extreme temperatures. The links to droughts or periods of extreme low flows in rivers are more complex. [Global Circulation](#)

[Models](#) (GCMs) – a collection of numerical models that provide a 3-D analysis of global climate interactions such as atmosphere, oceans, ice and land – [predict](#) increases in temperatures for some regions, such as southern Europe.

In terms of tropical cyclones, the effects of climate change on these phenomena is an active area of research as the processes are complex. For example, the [Clausius-Clapeyron](#) relationship which can be related to the water vapour-carrying capacity of the atmosphere, may have an impact on the strength and intensity of such storms.

The relationship states that for every degree rise in temperature, the water-holding capacity of the atmosphere increases by 7%, so in a warming ocean, the air above the water has a much greater capacity to hold water and thus store more rain that can feed more powerful storms.

But sinking cold air from the upper atmosphere may prevent storms from rising in the first place. If this happens more frequently with climate change then we can expect fewer such storms. That means in the future there may be fewer tropical cyclones forming, but those that do will be stronger and more intense.

Wet wet wet

In a warming world, we can expect it to get wetter. The distribution of the rainfall throughout the year could change as we experience longer, drier spells, although when rain falls it may be in intense bursts. Recent [research](#) by Newcastle University analysed the results from finer scale GCMs climate projections and suggests we may expect more intense summer rainfall in the UK in future. [New climate projections](#) from GCMs are being prepared for the UK to help predict what the future climate may look like.

Flooding from intense rainfall or river sources has many complex drivers which cause the damage in catchment areas. For example, land use changes (such as intensive farming practices or deforestation) and the degree of urbanisation both play a part in flood risk. Recent [research](#) for the UK suggests that we will see an increase in the frequency of extreme river flooding.

So what can we expect for 2018? Already Australia is experiencing [extreme heat](#), while the eastern seaboard of the US is suffering a [severe cold spell](#), and the west coast [devastating mudslides](#) that have killed 17 people. So far 2018 seems to be picking up right where 2017 left off.

Governments need to recognise and absorb that extreme weather across the globe is likely to become more common and start to adapt accordingly, rather than treat it as shocking one-off events. Otherwise we risk increasing loss of life and environmental damage in the future.

This article was originally published on [The Conversation](#). Read the [original article](#).

Provided by The Conversation

Citation: Will the weather get worse in 2018? What the experts say (2018, January 12) retrieved 20 September 2024 from <https://phys.org/news/2018-01-weather-worse-experts.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.