

Researchers explore anti-inflammatory activity of curcumin

17 January 2018



Credit: iStock

Extracts of the plant turmeric—the spice that gives Indian curries a yellow color—have been used as an anti-inflammatory treatment in traditional Asian medicine for centuries. Clinical trials of curcumin (the active chemical compound in turmeric), however, have produced mixed results. A molecular understanding of curcumin's biological effects is needed.

Claus Schneider, Ph.D., and colleagues have now discovered that [curcumin](#) is a "pro-drug" that is converted into reactive metabolites with anti-inflammatory activities. The metabolites of curcumin, produced by [oxidation reactions](#), covalently bind to and inhibit proteins in the inflammatory NF-kappa-B signaling pathway.

The researchers found that curcumin undergoes oxidation reactions readily in vitro. They suggest that insufficient bioactivation in vivo may explain the mixed results in human studies of curcumin activity.

The findings, reported in the *Journal of Biological Chemistry*, suggest that metabolic bioactivation

should be considered in [clinical trials](#) of curcumin and other dietary polyphenols of medicinal interest, such as resveratrol (red wine), quercetin (onions) and epigallocatechin gallate (green tea).

More information: Rebecca L. Edwards et al. The anti-inflammatory activity of curcumin is mediated by its oxidative metabolites, *Journal of Biological Chemistry* (2017). DOI: [10.1074/jbc.RA117.000123](https://doi.org/10.1074/jbc.RA117.000123)

Provided by Vanderbilt University

APA citation: Researchers explore anti-inflammatory activity of curcumin (2018, January 17) retrieved 29 September 2020 from <https://phys.org/news/2018-01-explore-anti-inflammatory-curcumin.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.