


Study helps explain Greenland glaciers' varied vulnerability to melting

14 March 2018




Photographs taken of the Mogens North glacier in southeast Greenland in 1932 and 2013 show how much ice has been lost in the past eight decades. UCI glaciologists have created new maps of this part of Greenland using data from NASA missions and learned why some of the massive, moving ice slabs are more vulnerable to melting than others. Credit: National History Museum of Denmark (left) and Hans Henrik Tholstrup

Using data from NASA missions observing Earth, researchers at the University of California, Irvine have created new maps of the bed topography beneath a score of glaciers in southeast Greenland, thereby gaining a much better understanding of why some are undergoing rapid retreat and others are relatively stable.

"The undersides of [glaciers](#) in deeper valleys are exposed to warm, salty Atlantic [water](#), while the others are perched on sills, protected from direct exposure to warmer [ocean](#) water," said Romain Millan, lead author of the study, available online in the American Geophysical Union journal *Geophysical Research Letters*. "We have been able to demonstrate unequivocally that glacier retreat in southeast Greenland is controlled by the topography of the bedrock under the ice and by ocean temperature."

Millan, a UCI graduate student researcher in Earth system science, and his colleagues analyzed 20 major outlet glaciers in southeast Greenland using high-resolution airborne gravity measurements and ice thickness data from NASA's Operation IceBridge mission; bathymetry information from NASA's Oceans Melting Greenland project; and results from the BedMachine version 3 computer model, developed at UCI.

They found glacial fjords hundreds of meters deeper than previously estimated; the full extent of the marine-based portions of the glaciers; deep troughs enabling Atlantic Ocean water to reach the glacier fronts and melt them from below; and few shallow sills that limit contact with this warmer water.



Credit: University of California, Irvine

"It's important to understand the physical processes controlling the retreat in order to improve projections of [sea level rise](#) from this region in a

warming climate," Millan said. "Until recently, we had little information on ocean temperature and water depth in these fjords to quantify these processes, so the interpretation of glacier evolution on a case-by-case basis was difficult."

Co-author Eric Rignot, UCI professor of Earth system science, added, "Now that the picture is clear, the role of the ocean in glacier evolution is overwhelming."

Rignot, who has led dozens of research expeditions to Earth's polar regions, said that southeast Greenland - with its fast-moving glaciers, deep fjords and harsh climate conditions - poses significant challenges to researchers.

"Thanks to the newest NASA missions, such as Oceans Melting Greenland and Operation IceBridge, we have been able to make great advances in understanding the evolution of this very dynamic sector of Greenland and its impact on sea level rise now and in decades to come," he said.

More information: R. Millan et al. Vulnerability of Southeast Greenland glaciers to warm Atlantic Water from Operation IceBridge and Ocean Melting Greenland data., *Geophysical Research Letters* (2018). [DOI: 10.1002/2017GL076561](https://doi.org/10.1002/2017GL076561)

Provided by University of California, Irvine
APA citation: Study helps explain Greenland glaciers' varied vulnerability to melting (2018, March 14) retrieved 24 June 2021 from <https://phys.org/news/2018-03-greenland-glaciers-varied-vulnerability.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.