

# Marker for cancer stem cells—Fluorescent probe identifies tumor-initiating cells

4 April 2018

Cells with stem-cell characteristics appear to be especially important in the formation and metastasis of tumors. Scientists have now developed a universal fluorescent probe for these "cancer stem cells". As reported in the journal *Angewandte Chemie*, this dye also selectively kills off the cells.

"Cancer stem cells", also known as [tumor](#)-initiating cells (TIC), appear to cause relapses after radiation and chemotherapy because a single surviving TIC can cause a new tumor to grow. In addition, they appear to be the main cause of metastasis. Effective tumor treatment must therefore aim to kill off TICs as extensively as possible. To this end, a "probe" that marks these [cancer](#) stem cells would be useful so that they become visible. Although there are markers that also recognize TICs associated with some types of cancer, no universal, selective probe for [cancer stem cells](#) has been found.

A team from the Agency for Science Technology and Research (A\*STAR) in Singapore, Pohang University of Science and Technology in Korea, and other research organizations in Singapore and Korea, have now succeeded in finding such a probe. They were able to show that their new probe, a fluorescent dye, selectively stains TICs from a broad variety of cancers, including tumors of the lung, central nervous system, breast, kidney, ovary, colon, and prostate, as well as melanomas. Healthy cells and "ordinary" tumor cells were not marked. At high concentrations, the dye also demonstrates considerable cytotoxicity toward TIC, while other cells are barely affected.

Led by Nam-Young Kang and Young-Tae Chang, the researchers discovered that their probe, named TiY (for tumor-initiating cell probe yellow), recognizes vimentin, which is a molecule in the cytoskeleton. Vimentin is more concentrated in [epithelial cells](#) when they transform into mesenchymal cells. Epithelial cells form the tissue

that covers the inner and outer surfaces of the body, forming a boundary with the environment. The cells are polar, meaning that the side facing toward the underlying tissue and the side directed outward toward the lumen are different. The [cells](#) are also firmly integrated into the cell wall. When they transform into [mesenchymal cells](#), they lose their polarity, are freed from the cell structure, and may wander. This process plays an important role in the development of embryos and healing of wounds. It is also involved in the metastasis of tumors.

As the first fluorescent probe for TIC, TiY could be a valuable tool for the visualization and isolation of TIC, and may help in the development of tumor treatments that are targeted toward zones concentrated with vimentin, thereby inhibiting relapses.

**More information:** Yong-An Lee et al, Identification of Tumor Initiating Cells with a Small-Molecule Fluorescent Probe by Using Vimentin as a Biomarker, *Angewandte Chemie International Edition* (2018). DOI: [10.1002/anie.201712920](https://doi.org/10.1002/anie.201712920)

Provided by Wiley

APA citation: Marker for cancer stem cells—Fluorescent probe identifies tumor-initiating cells (2018, April 4) retrieved 3 March 2021 from <https://phys.org/news/2018-04-marker-cancer-stem-cellsfluorescent-probe.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*