

# Climate change impacts fragile river ecosystems

May 16 2018

---



South Africa's Kruger National Park. Credit: Stephen Tooth

Boulder, Colo., USA: Research undertaken in South Africa's Kruger National Park (KNP) has shown that some of the world's most sensitive and valuable riverine habitats are being destroyed due to an increasing

frequency of cyclone-driven extreme floods.

As part of a Natural Environmental Research Council (UK) funded project, researchers from the universities of Hull, Aberystwyth, and Salford and the engineering consultants "Architecture, Engineering, Consulting, Operations, and Maintenance" (AECOM), used laser survey technology (LiDAR) flown from an aircraft, to measure the impacts of cyclone-driven extreme floods in 2000 and 2012 on rivers in the KNP. The KNP game reserve has global significance for its habitats and associated species, and the rivers flowing through the park provide essential ecosystem services, including water and [habitat](#) in the shape of the many varied channel morphologies and associated riparian forest. The high-resolution data has been used to create accurate digital models of the [river bed](#), and through comparisons with pre-2012 flood data, they were able map detailed spatial patterns of erosion and deposition.

Dr. David Milan, University of Hull, principal investigator for the project said, "We are primarily interested in trying to understand how these large bedrock-influenced river channels respond to large floods. From comparing our LiDAR models between 2012 and 2004, we have calculated that the 2012 event alone removed almost 1.25 million tonnes of sediment from the river bed. We also found that patches of mature riparian forest that survived larger floods in 2000 were removed by the 2012 floods. There is a suggestion that the frequency of large flood events is increasing due to climate change, and our analysis of river channel morphology for a 50 km length of the Sabie River shows us that these rivers need timespans longer than a decade to recover."

The UK group has concerns over the impacts of the geomorphological changes happening in the KNP rivers. Drylands are known to be some of the regions most vulnerable to climate change, and some climate model predictions suggest increased landfall for cyclones over South Africa.

Dr. Milan continued, "We present a conceptual model showing the likely pathways that the KNP river systems are likely to follow in the future depending upon [flood](#) frequency and magnitude, and conclude by suggesting that more frequent floods will continue to strip out sediment and vegetation from the river channel, leaving a more barren environment with less habitat value. Continued progressive loss of habitat diversity will fundamentally, and for all intents and purposes irreversibly, alter our riverine landscapes and this will be accompanied by a catastrophic loss of species unable to adapt to the new environments. Conservationists need to work alongside geomorphologists to look at ways in which dryland river habitats can be best managed into the future."

These results not only have significance for the rivers of the KNP, but also for bedrock-influenced rivers in other dryland areas globally.

The full findings of this research have just been published in the *Geological Society of America Bulletin*.

**More information:** David Milan et al. Morphodynamics of bedrock-influenced dryland rivers during extreme floods: Insights from the Kruger National Park, South Africa, *GSA Bulletin* (2018). [DOI: 10.1130/B31839.1](#)

Provided by Geological Society of America

Citation: Climate change impacts fragile river ecosystems (2018, May 16) retrieved 21 September 2024 from <https://phys.org/news/2018-05-climate-impacts-fragile-river-ecosystems.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.