

# Newly discovered protein operated in earliest organisms

17 May 2018



Microscopic image of cells of the cyanobacterium *Synechocystis* sp. PCC6803. Photo: Khaled Selim. Credit: Universitaet Tübingen

Life on Earth depends on photosynthetic carbon dioxide (CO<sub>2</sub>) fixation to form organic carbon. Plants take atmospheric CO<sub>2</sub> and transform it into organic molecules such as glucose. This process evolved in cyanobacteria and was later conveyed to eukaryotes, giving rise to plastids in algae and plants. Researchers have now discovered a new protein which is involved in this complex process; it regulates the intake of CO<sub>2</sub> into the cell. Khaled Selim and Professor Karl Forchhammer from the Interfaculty Institute for Microbiology and Infection Medicine (IMIT), together with colleagues from the Max-Planck Institute for Proteinevolution and from the University Rostock, describe their finding of the conserved cyclic AMP receptor protein, SbtB, in the latest edition of *Proceedings of the National Academy of Sciences (PNAS)*.

SbtB participates in the sensing of fluctuating ambient CO<sub>2</sub> concentrations to adjust CO<sub>2</sub> fixation to different environments. SbtB represents a new member of the PII signal transduction superfamily, known for binding the energy carrier molecules ATP and ADP - a kind of battery for the cell. The newly discovered protein, however, also binds the

cyclic nucleotide cAMP, which plays fundamental roles in all organisms for signalling the state of carbon metabolism. Up to now, cAMP was chiefly known for being a key messenger molecule, required in the maintenance of the organisms glucose balance - used for instance in regulating blood-sugar levels.

SbtB is the first protein known to bind cAMP to regulate the CO<sub>2</sub> metabolism in cyanobacteria. In identifying SbtB, the researchers have found new principle of carbon sensing through cAMP, which is important for acclimation to varying C<sub>i</sub> regimes in the ecological niches of [cyanobacteria](#). Cyanobacteria, also known as blue-green algae, are among the oldest group of organisms on Earth; knowing more about their workings offers clues to very early life on the planet.

**More information:** PII-like signaling protein SbtB links cAMP sensing with cyanobacterial inorganic carbon response. *Proceedings of the National Academy of Sciences (PNAS)*, [doi.org/10.1073/pnas.1803790115](https://doi.org/10.1073/pnas.1803790115)

Provided by Universitaet Tübingen

APA citation: Newly discovered protein operated in earliest organisms (2018, May 17) retrieved 24 October 2021 from <https://phys.org/news/2018-05-newly-protein-earliest.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*