

Plant symbioses—fragile partnerships

25 May 2018



Credit: CC0 Public Domain

All plants require an adequate supply of inorganic nutrients, such as fixed nitrogen (usually in the form of ammonia or nitrate), for growth. A special group of flowering plants thus depends on close symbiotic relationships with nitrogen-fixing bacteria as a source of fixed nitrogen. The plant provides specialized root nodules as housing for its bacterial benefactors, which allows them to make use of energy-rich organic carbon compounds in return for the ammonia they generate from atmospheric nitrogen. Thus, both partners reap advantages from the relationship.

Nevertheless, a comparative study of 37 [plant genomes](#) carried out by Munich molecular biologists based at Ludwig-Maximilians-University Munich (LMU) and the Helmholtz Zentrum München (HMGU) in cooperation with French and Chinese colleagues, demonstrates that the capacity to form root-nodule partnerships has repeatedly been lost during the course of evolution. The new findings appear in the leading journal *Science*.

The ability to form root nodules in response to the

presence of nitrogen-fixing bacteria is currently restricted to four orders of flowering [plants](#) (Fabales, Fagales, Cucurbitales and Rosales), to which many agriculturally significant species belong, including beans, peas and soya. The authors of the new study therefore assumed that the genetic basis for this type of symbiosis initially evolved in a common ancestor of these groups.

"However, only 10 of the 28 plant families that make up these four orders are now able to form nitrogen-fixing root nodules. Moreover, most of the genera in nine of these families do not adopt the symbiotic lifestyle," says LMU geneticist Professor Martin Parniske, who initiated the international collaboration. In order to identify the genetic grounds for the relatively low incidence of root nodule symbioses among the [plant families](#) in which it occurs, the consortium collected DNA samples and the BGI in China sequenced the entire genome of 10 species that form distinct types of nodules with diverse bacterial symbionts, which employ different infection strategies. In close cooperation with LMU and the HMGU, the genomic sequence data for these species were then included in a genome-wide comparison of 37 plant species representing the four orders mentioned above. Much to everyone's surprise, the analyses revealed that many members of these groups—including well known cultivars such as strawberries, blackberries and apples—had once possessed, but subsequently lost the capacity to form symbioses.

"We discovered that the gene NODULE INCEPTION (NIN), which is an essential prerequisite for root nodule symbiosis, has been repeatedly and independently disabled by mutation in several different lineages. The functional gene was therefore most probably present in a [common ancestor](#) of these lines, but was later lost several times independently," explains first author Maximilian Griesmann, doctoral student at LMU who at HMGU worked closely with Manuel Spannagl, Georg Haberer and Klaus F.X. Mayer, head of the Department of Genomics and Systems

Biology of Plant Genomes at HMGU, on the genetic analyses. "These findings were completely unexpected. In many terrestrial ecosystems, low nitrogen availability is a factor limiting plant growth, so that one would expect the symbiosis to be associated with a definite selective advantage," adds Parniske.

According to the paper's authors, one possible reason for the frequent breakdown of these symbioses is that root nodules can be invaded by parasitic bacteria, which compete for and consume the compounds provided by the plants without supplying nitrogen in return. In that case, loss of the symbiosis should effectively be favored as an evolutionary adaptation that enables the plant to escape this form of exploitation. Other scenarios in which the symbiosis no longer benefits the host are also conceivable, such as a persistent oversupply of nitrogen or situations in which the availability of other essential substances like water or phosphates constrains plant growth.

In the field of synthetic biology, efforts to equip non-symbiotic crop plants with the ability to establish root-nodule symbioses are underway, with a view to reducing the need to supply fixed nitrogen by applying environmentally problematic fertilizers. "Our results suggest that evolution exerts a selective pressure against the maintenance of root-nodules symbioses, and future studies need to take this factor into account," Mayer says.

More information: "Phylogenomics reveals multiple losses of nitrogen-fixing root nodule symbiosis" *Science* (2018).

science.sciencemag.org/lookup/.../1126/science.aat1743

Provided by Ludwig Maximilian University of Munich

APA citation: Plant symbioses—fragile partnerships (2018, May 25) retrieved 25 January 2021 from <https://phys.org/news/2018-05-symbiosesfragile-partnerships.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.