


# The true power of the solar wind

12 June 2018



Extremely fast particles from the sun hit Mercury. Credit: NASA, montage: TU Wien

Friedrich Aumayr from the Institute of Applied Physics at TU Wien. These particles hit the [surface](#) rocks at a speed of 400 to 800 km per second and the impact can eject numerous other atoms. These particles can rise high before they fall back to the surface, creating an "exosphere" around the Moon or Mercury—an extremely thin atmosphere of atoms sputtered from the surface rocks by solar wind bombardment.


This exosphere is of great interest for space research because its composition allows scientists to deduce the chemical composition of the [rock](#) surface—and it is much easier to analyse the exosphere than to land a spacecraft on the surface. In October 2018, ESA will send the BepiColombo probe to Mercury, which is to obtain information about the geological and chemical properties of Mercury from the composition of the exosphere.

The planets and moons of our solar system are continuously being bombarded by particles hurled away from the sun. On Earth this has hardly any effect, apart from the fascinating northern lights, because the dense atmosphere and the magnetic field of the Earth protect us from these solar wind particles. But on the Moon or on Mercury things are different: There, the uppermost layer of rock is gradually eroded by the impact of sun particles.

New results of the TU Wien now show that previous models of this process are incomplete. The effects of solar wind bombardment are in some cases much more drastic than previously thought. These findings are important for the ESA mission BepiColombo, Europe's first Mercury mission. The results have now been published in the planetology journal *Icarus*.

## An exosphere of shattered rock

"The solar wind consists of charged [particles](#)—mainly hydrogen and helium ions, but heavier atoms up to iron also play a role," explains Prof.



Paul Szabo in the lab at TU Wien. Credit: Vienna University of Technology

## Charge matters

However, this requires a precise understanding of the effects of the solar wind on the rock surfaces,


and this is precisely where decisive gaps in knowledge still exist. Therefore, the TU Wien investigated the effect of ion bombardment on wollastonite, a typical moon rock. "Up to now it was assumed that the kinetic energy of the fast particles is primarily responsible for atomization of the rock surface," says Paul Szabo, Ph.D. student in Friedrich Aumayr's team and first author of the current publication. "But this is only half the truth: we were able to show that the high electrical charge of the particles plays a decisive role. It is the reason that the particles on the surface can do much more damage than previously thought."

When the particles of the solar wind are multiply charged, i.e. when they lack several electrons, they carry a large amount of energy which is released in a flash on impact. "If this is not taken into account, the effects of the solar [wind](#) on various rocks are misjudged," says Paul Szabo. Therefore, it is not possible to draw exact conclusions about the surface rocks with an incorrect model from the composition of the exosphere.

positively. And the contribution of heavier ions with an even greater [electrical charge](#) must not be neglected either.

A cooperation of different research groups was necessary for these findings: High-precision measurements were carried out with a specifically developed microbalance at the Institute of Applied Physics. At the Vienna Scientific Cluster VSC-3 complex computer simulations with codes developed for nuclear fusion research were carried out in order to be able to interpret the results correctly. The Analytical Instrumentation Center and the Institute for Chemical Technologies and Analytics of the TU Vienna also made important contributions.

**More information:** Paul S. Szabo et al. Solar Wind Sputtering of Wollastonite as a Lunar Analogue Material – Comparisons between Experiments and Simulation, *Icarus* (2018). [DOI: 10.1016/j.icarus.2018.05.028](https://doi.org/10.1016/j.icarus.2018.05.028)



Provided by Vienna University of Technology

Herbert Biber, Paul Szabo, Friedrich Aumayr, Reinhard Stadlmayr, Daniel Mayer (left to right). Credit: Vienna University of Technology

Protons make up by far the largest part of the [solar wind](#), and so it was previously thought that they had the strongest influence on the rock. But as it turns out, helium actually plays the main role because, unlike protons, it can be charged twice as

APA citation: The true power of the solar wind (2018, June 12) retrieved 22 September 2019 from <https://phys.org/news/2018-06-true-power-solar.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*