

# Neonics are being ingested by free-ranging animals, study finds

20 June 2018

Health impacts of neonicotinoids may go well beyond bees, according to a new University of Guelph study.

U of G researchers found residues of the insecticides in the livers of [wild turkeys](#), providing evidence that this common agrochemical is being ingested by free-ranging animals.

The researchers from the Ontario Veterinary College are among the first to study the broader effects of neonics on wildlife.

Published in *Environmental Science and Pollution Research*, the study showed that nearly 10 of the 40 wild turkey carcasses tested had detectable levels of neonicotinoids in their livers. Two types of the insecticide were found in some birds.

The researchers also found corn and soybean seeds coated with the insecticide in the digestive system of some birds.

"Wild turkeys supplement their diet with seeds from farm fields," said pathobiology professor Claire Jardine. She conducted the study with former pathobiology professor Nicole Nemeth, who is now at the University of Georgia, pathobiology Ph.D. Amanda MacDonald, and Philippe Thomas, a wildlife toxicologist with Environment and Climate Change Canada.

"There has been growing concern among natural resource managers, conservationists and hunters about whether the use of neonics may be linked to poor reproductive output of wild turkeys."

While researchers have focused on health risks of neonicotinoids to bees, studying exposure levels in larger wildlife species is critical in understanding wider impacts on migratory behaviour, reproduction and mortality, said Jardine.

"Our results serve as baseline data for southern

Ontario wild turkeys and provide context for reference values in future analyses."

MacDonald began the study after officials with the Ontario Federation of Hunters and Anglers called for research into the potential threat posed by neonics to wild turkeys.

"A number of member hunters throughout southern Ontario had seen wild turkeys in the fields eating these seeds," said MacDonald. "In certain areas, they noticed a lack of young birds and wanted to know if neonicotinoids had anything to do with it."

The study proves wild [turkeys](#) consumed neonic-treated seeds, but long-term health effects on the birds remain unknown, added MacDonald.

Previous studies have found that neonic-coated seeds cause [health risks](#) in partridges, pigeons and quail. Small amounts of the insecticide have been shown to affect body mass, reproductive efforts and perhaps mortality in migratory white-crowned sparrows.

"We need to continue to assess levels of neonics in a variety of wildlife, especially those that may feed off the ground or consume plants and insects and therefore might be more likely to come into contact with them," said Nemeth.

**More information:** Amanda M. MacDonald et al, Neonicotinoid detection in wild turkeys (*Meleagris gallopavo silvestris*) in Ontario, Canada, *Environmental Science and Pollution Research* (2018). [DOI: 10.1007/s11356-018-2093-0](https://doi.org/10.1007/s11356-018-2093-0)

Provided by University of Guelph

APA citation: Neonics are being ingested by free-ranging animals, study finds (2018, June 20) retrieved 18 April 2021 from <https://phys.org/news/2018-06-neonics-ingested-free-ranging-animals.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*