

Understanding the universe through neutrinos

16 July 2018, by Anne M Stark



than 1016 times the age of the universe and could help determine whether neutrinos are their own antiparticles and explain why, from equal parts of matter and antimatter, the universe evolved into its current matter-dominated state.


The design of the nEXO detector—a 5-ton liquid xenon (Xe) time projection chamber (TPC) using 90 percent enriched ^{136}Xe —takes advantage the best technology for the next phase of NDBD search.

The xenon vessel and vacuum vessel for the next Enriched Xenon Observatory (nEXO) experiment was built at Lawrence Livermore National Laboratory. The experiment will search for an extremely rare nuclear process called neutrinoless double-beta decay (NDBD). Credit: Lawrence Livermore National Laboratory

Determining features of the elusive particle known as a neutrino – through the observation of an extremely rare nuclear process called neutrinoless double-beta decay (NDBD)—could provide a glimpse into the nature of the universe during the earliest moments of the Big Bang.

As part of an international collaboration, Lawrence Livermore National Laboratory (LLNL) scientists have proposed the next Enriched Xenon Observatory (nEXO) experiment, a candidate for the next generation of NDBD experiments. If discovered, NDBD would demonstrate the existence of a new elementary particle, the Majorana fermion. This discovery could reshape the standard model of particle physics and lead to a better understanding of neutrinos and their impact on the evolution of the [universe](#). The research behind the experiment appears in the journal *Physical Review C*.

NDBD is a theoretical process with a half-life more



The Enriched Xenon Observatory 200 (EXO-200) experiment provides the basis for current work on a more sensitive detector for observing neutrinoless double beta decay (NDBD). Shown here are the EXO-200 readout wires and avalanche photodiodes used to measure induced and collected charge and scintillation light from particle decays in the detector's main vessel. Credit: SLAC National Accelerator Laboratory

"A competitive 2-order-of-magnitude increase in NDBD half-life sensitivity over current experiments is possible" using the nEXO detector, said LLNL scientist Samuele Sangiorgio, lead author of the paper. "We now have great confidence in nEXO's

design and approach, and we will be able to measure this rare event."

Scientists expect to see only about a dozen decays in a decade-long experiment. Because of this very low signal rate, false signals from background radiation and cosmic rays must be suppressed as much as is feasible. "Understanding the backgrounds is key to a make a convincing case for a NDBD experiment, and indeed is one of the main aspects of the paper," Sangiorgio said.

More information: J. B. Albert et al. Sensitivity and discovery potential of the proposed nEXO experiment to neutrinoless double- β decay, *Physical Review C* (2018). [DOI: 10.1103/PhysRevC.97.065503](https://doi.org/10.1103/PhysRevC.97.065503)

Provided by Lawrence Livermore National Laboratory

APA citation: Understanding the universe through neutrinos (2018, July 16) retrieved 26 June 2019 from <https://phys.org/news/2018-07-universe-neutrinos.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.